

# Around the Approximation Property for Quantum Groups

Matthew Daws

UCLan

Quantum Groups Seminar, January 2023

## Fejér's Theorem

Let's recall this from classical Fourier Analysis. Identify  $\mathbb{T} = [-\pi, \pi)$  which has Haar measure  $\frac{ds}{2\pi}$ . For a “nice” function  $f$  on  $\mathbb{T}$  define

$$c_k = \int_{-\pi}^{\pi} f(s) e^{-iks} \frac{ds}{2\pi}, \quad s_n(f, x) = \sum_{k=-n}^n c_k e^{ikx}.$$

### Theorem

For  $f \in C(\mathbb{T})$ , the Cesàro sums

$$\sigma_n(f, x) = \frac{1}{n} \sum_{k=0}^{n-1} s_k(f, x)$$

converge uniformly to  $f(x)$ .

## Think about this in a “quantum” framework

For me, the Fourier transform is between Hilbert spaces:

$$\mathcal{F} : L^2(\mathbb{T}) \rightarrow \ell^2(\mathbb{Z}); \quad f \mapsto (c_k) = \left( \int_{-\pi}^{\pi} f(s) e^{-iks} \frac{ds}{2\pi} \right).$$

Let  $C(\mathbb{T})$  act on  $L^2(\mathbb{T})$ . For  $n \in \mathbb{Z}$ , let  $\lambda_n$  be the translation operator on  $\ell^2(\mathbb{Z})$ , and let  $C_r^*(\mathbb{Z})$  be the closed linear span of such operators.

We then obtain

$$\mathcal{F}_0 : C(\mathbb{T}) \rightarrow C_r^*(\mathbb{Z}); \quad f \mapsto \mathcal{F}f\mathcal{F}^{-1}.$$

Calculation shows:

$$\mathcal{F}_0^{-1} : \lambda_n \mapsto (e^{ins})_{s \in \mathbb{T}}.$$

## Continued 1: Normal functionals

For a  $C^*$ -algebra  $A \subseteq \mathcal{B}(H)$ , given  $\xi, \eta \in H$ , let  $\omega_{\xi, \eta} \in A^*$  be the (normal) functional

$$A \ni a \mapsto (a\xi|\eta) \in \mathbb{C}.$$

Think of  $L^1(\mathbb{T})$  as those functionals on  $C(\mathbb{T})$  of this form. Indeed, given  $\xi, \eta \in L^2(\mathbb{T})$  and  $f \in C(\mathbb{T})$ ,

$$\langle f, \omega_{\xi, \eta} \rangle = (f\xi|\eta) = \int_{-\pi}^{\pi} f(s)\xi(s)\overline{\eta(s)} \frac{ds}{2\pi} = \langle f, \xi\bar{\eta} \rangle.$$

Similarly, the *Fourier Algebra*  $A(\mathbb{Z})$  is the collection of such normal functionals on  $C_r^*(\mathbb{Z})$ . (That this is a *closed subspace* is true, but not obvious).

## Continued 2: Function Spaces

Given  $\omega = \omega_{\xi, \eta} \in A(\mathbb{Z})$ , we can identify this with a function on  $\mathbb{Z}$  by

$$\omega \leftrightarrow (\omega(n))_{n \in \mathbb{Z}} = (\langle \lambda_{-n}, \omega \rangle)_{n \in \mathbb{Z}}.$$

As  $C_r^*(\mathbb{Z})$  is the span of  $\{\lambda_n : n \in \mathbb{Z}\}$ , the values  $\{\omega(n) : n \in \mathbb{Z}\}$  determines  $\omega$ . Use of “ $-n$ ” seems odd, but makes things work (and occurs in the general quantum theory).

Recall  $\mathcal{F}_0 : C(\mathbb{T}) \rightarrow C_r^*(\mathbb{Z})$ . The Banach space adjoint restricts to give

$$\mathcal{F}_1 = \mathcal{F}_0^* : A(\mathbb{Z}) \rightarrow L^1(\mathbb{T}); \omega_{\xi, \eta} \mapsto \omega_{\mathcal{F}^*(\xi), \mathcal{F}^*(\eta)}.$$

This is a bijection, and the inverse  $L^1(\mathbb{T}) \rightarrow A(\mathbb{Z})$  is just the usual Fourier transform (thought of as acting between function spaces).

## Continued 3: Algebras

$L^1(\mathbb{T})$  is an algebra under convolution, and  $A(\mathbb{Z})$  is an algebra of functions with the pointwise product.

- $\mathcal{F}_1$  is a homomorphism.
- $\mathcal{F}_0$  is a homomorphism.

$A(\mathbb{Z})$  acts on its dual space, and this restricts to turn  $C_r^*(\mathbb{Z})$  into an  $A(\mathbb{Z})$ -module. Similarly for  $L^1(\mathbb{T})$ ,

$$\omega \cdot \lambda_n = \omega(n)\lambda_n, \quad f \cdot F = F \star \check{f} \quad (F \in C(\mathbb{T}), f \in L^1(\mathbb{T})).$$

Here  $\check{f}(s) = f(-s)$ .

- $\mathcal{F}_0$  is a module homomorphism.

## Back to Fejér

For  $F \in C(\mathbb{T})$  we have

$$\sigma_n(F, \cdot) = F \star F_n = \check{F}_n \cdot F,$$

where  $F_n \in L^1(\mathbb{T})$  is the Fejér kernel; we have  $\check{F}_n = F_n$ .

Push this through  $\mathcal{F}_0$  to obtain  $\omega_n = \mathcal{F}_1(\check{F}_n)$  with

$$\omega_n \cdot a = \mathcal{F}_0(F_n \cdot F_0^{-1}(a)) \xrightarrow{n \rightarrow \infty} \mathcal{F}_0(F_0^{-1}(a)) = a \quad (a \in C_r^*(\mathbb{Z})).$$

Indeed,  $\omega_n$ , as a function on  $\mathbb{Z}$ , is simply the “triangle”, piecewise linear with  $\omega_n(0) = 1$  and  $\omega_n(n) = \omega_n(-n) = 0$ .

*We obtain a sequence of (normalised, positive definite) functions in  $A(\mathbb{Z})$  which acts on  $C_r^*(\mathbb{Z})$  as an “approximate identity”.*

We can replace  $\mathbb{Z}$  by an arbitrary group; and/or weaken some conditions.

## Amenability

Let  $G$  be a locally compact (or just discrete) group. Form  $L^2(G)$ , form the translation operators  $\{\lambda_g : g \in G\}$ . (When  $G$  is discrete...) The norm closed linear span is  $C_r^*(G)$ , and the bicommutant is  $VN(G)$ . The predual of  $VN(G)$  is  $A(G)$ , the *Fourier Algebra*, considered as an algebra of functions in the same way,  $\omega \leftrightarrow (\omega_g)_{g \in G} = (\langle \lambda_{g^{-1}}, \omega \rangle)_{g \in G}$ . Turn  $C_r^*(G)$  and  $VN(G)$  into  $A(G)$ -modules for the dual action.

### Theorem

*The following are equivalent:*

- *$A(G)$  contains a net of normalised positive definite functions (i.e. normal states on  $VN(G)$ ) which form an approximate identity for  $C_r^*(G)$ , or a weak\*-approximate identity for  $VN(G)$ ;*
- *$A(G)$  contains some bounded approximate identity;*
- *$G$  is amenable.*

# A-T-menability or the Haagerup property

## Question

Can we expand the space of functions away from  $A(G)$  to obtain a larger class of groups than those which are amenable?

Instead of using the predual of  $VN(G)$ , could we use the dual of  $C_r^*(G)$ ? No: if this has a bai then it has a unit, and  $G$  is amenable. Could we use the dual of the full group  $C^*$ -algebra  $C^*(G)$ ? No: this is always unital. But all functions in  $A(G)$  vanish at infinity.

## Definition

$G$  has the *Haagerup Property* if there is a net of normalised positive-definite functions which vanish at infinity, and converge to 1 uniformly on compacta.

## Completely bounded multipliers

A key property of  $A(G)$  functions is that they “multiply” (or act on)  $C_r^*(G)$  and  $VN(G)$ . Of course,  $A(G)$  is an algebra.

### Definition

A *multiplier* of  $A(G)$  is a function  $f$  on  $G$  such that  $f\omega \in A(G)$  for each  $\omega \in A(G)$ .

Such an  $f$  is automatically continuous, and by the Closed Graph Theorem, the resulting map  $A(G) \rightarrow A(G); \omega \mapsto f\omega$  is continuous.  $f$  acts on  $VN(G)$  and by restriction, on  $C_r^*(G)$ .

### Definition

A multiplier  $f$  is *completely bounded* if the resulting map on  $VN(G)$ , say  $M_f$ , (equivalently  $C_r^*(G)$ ) is completely bounded.

$$M_f \otimes \text{id} : VN(G) \otimes \mathbb{M}_n \rightarrow VN(G) \otimes \mathbb{M}_n.$$

## Weak amenability

Of course, each  $\omega \in A(G)$  is itself a (cb-)multiplier.

### Theorem (Losert)

*The following are equivalent:*

- $G$  is amenable
- the map from  $A(G)$  into the algebra of multipliers of  $A(G)$  is bounded below;
- the map from  $A(G)$  into the algebra of cb-multipliers of  $A(G)$  is bounded below.

### Definition

$G$  is *weakly amenable* if there is a net  $(\omega_i)$  in  $A(G)$ , bounded in the  $\|\cdot\|_{cb}$  norm, forming an approximate identity for  $C_r^*(G)$ .

E.g. (Haagerup)  $\mathbb{F}_2$ .

## The approximation property

The space of cb-multipliers,  $M_{cb}A(G)$ , is a dual space (and a dual Banach algebra).

- Each  $f \in L^1(G)$  defines a bounded functional on  $M_{cb}A(G)$  (by integration of functions).
- The closure of such functionals in  $M_{cb}A(G)^*$ , say  $Q_{cb}A(G)$ , is a predual for  $M_{cb}A(G)$ .

### Definition

$G$  has the *approximation property* (AP) when there is a net  $(\omega_i)$  in  $A(G)$  which converges to 1 weak\* in  $M_{cb}A(G)$ .

If such a net is bounded in  $M_{cb}A(G)$  then  $G$  is already weakly amenable.

## Examples

The class of groups with the AP is closed under extensions, while the class of weakly amenable groups is not (not even closed under semi-direct products).

- Let  $\Lambda_{cb}(G)$  be the infimum of  $M$  such that  $A(G)$  contains a net  $(\omega_i)$  converging to 1 on compacta, with  $\|\omega_i\|_{cb} \leq M$ .
- So  $G$  is weakly amenable exactly when  $\Lambda_{cb}(G) < \infty$ .
- [Cowling–Haagerup]  $\Lambda_{cb}(G_1 \times G_2) = \Lambda_{cb}(G_1)\Lambda_{cb}(G_2)$ .
- So if there is  $G$  with  $\Lambda_{cb}(G) > 1$  we can take an infinite product. Wreath products give such examples.
- This allows to find examples with AP which are not weakly amenable.
- In fact, much is known now about Lie groups and lattices therein.
- [Lafforgue–de la Salle]  $SL_3(\mathbb{Z})$  does not have the AP.

## Applications: finite-rank approximations

For those familiar with the notion of *nuclearity* the following should look slightly familiar.

### Definition

A  $C^*$ -algebra  $A$  has the *operator approximation property* (OAP) if there is a net of continuous finite-rank operators  $(\varphi_i)$  which converges to  $1_A$  in the point-stable topology:  $(\varphi_i \otimes \text{id})(u) \rightarrow u$  in norm, for each  $u \in A \otimes \mathcal{K}(\ell^2)$ .

### Theorem (Haagerup–Kraus)

For a discrete group  $G$  the following are equivalent:

- $G$  has the AP;
- $C_r^*(G)$  has the OAP.

Similar definitions/results hold for von Neumann algebras, and  $VN(G)$ .

## $L^p$ variants

We can replace  $L^2(G)$  by  $L^p(G)$  when defining the Fourier algebra and  $VN(G)$ . The operators  $(\lambda_s)_{s \in G}$  act on  $L^p(G)$  (by left-invariance of the Haar measure). The weak\*-linear span in  $\mathcal{B}(L^p(G))$  is  $PM_p(G)$ , the algebra of  $p$ -pseudo measures. Its predual is  $A_p(G)$  the *Figa-Talamanca–Herz algebra*.

We can also look at right-translation variants, leading to  $PM_p^r(G)$ . Let the commutant of this be  $CV_p(G)$ , the algebra of  $p$ -convolvers. We always have that  $CV_p(G) \supseteq PM_p(G)$ .

### Question

Is it true that  $CV_p(G) = PM_p(G)$ ?

## $L^p$ variants, continued

### Question

Is it true that  $CV_p(G) = PM_p(G)$ ?

Yes, if  $p = 2$ .

### Theorem (Cowling; see D.-Spronk)

*If  $G$  has the AP then  $CV_p(G) = PM_p(G)$*

The idea of the proof is that the net  $(\omega_i)$  in  $A(G)$  approximating the identity can be made to act on  $CV_p(G)$  in a way which weak\*-approximates the identity and which maps  $CV_p(G)$  into  $PM_p(G)$ .

# Locally compact quantum groups

(We assume the audience has some background).

Abstract object  $\mathbb{G}$  with:

- von Neumann algebra  $L^\infty(\mathbb{G})$  and  $C^*$ -algebra  $C_0(\mathbb{G})$ ;
- equipped with a coproduct  $\Delta : L^\infty(\mathbb{G}) \rightarrow L^\infty(\mathbb{G}) \bar{\otimes} L^\infty(\mathbb{G})$ ;
- which carry weights  $\varphi, \psi$  which are left/right invariant,

$$\varphi((\omega \otimes \text{id})\Delta(x)) = \varphi(x)\omega(1) \quad (x \in \mathcal{M}_\varphi^+, \omega \in L^1(\mathbb{G})^+).$$

From this, one gets:

- $L^1(\mathbb{G})$  becomes a Banach algebra, product induced by  $\Delta$ ;
- GNS for  $\varphi$  gives  $L^2(\mathbb{G})$  with  $L^\infty(\mathbb{G})$  in standard position;
- a multiplicative unitary  $W$ , so  $W_{12} W_{13} W_{23} = W_{23} W_{12}$ ;
- $C_0(\mathbb{G})$  is the closure of  $\{(\text{id} \otimes \omega)(W) : \omega \in \mathcal{B}(L^2(\mathbb{G}))_*\}$ .

# Duality

$$\lambda : L^1(\mathbb{G}) \rightarrow \mathcal{B}(L^2(\mathbb{G})); \quad \omega \mapsto (\omega \otimes \text{id})(W)$$

is a homomorphism. The closure of its image is a  $C^*$ -algebra  $C_0(\widehat{\mathbb{G}})$ .

- There indeed exists  $\widehat{\mathbb{G}}$  a LCQG;  $L^\infty(\widehat{\mathbb{G}})$  is the WOT closure.
- There is  $\widehat{\varphi}$  so that  $L^2(\widehat{\mathbb{G}}) = L^2(\mathbb{G})$  canonically.
- $W \in L^\infty(\mathbb{G}) \bar{\otimes} L^\infty(\widehat{\mathbb{G}})$  and  $\widehat{W} = \sigma(W^*)$  where  $\sigma$  is the swap map.

For  $G$  a locally compact group, we set  $L^\infty(\mathbb{G}) = L^\infty(G)$  with

$$\Delta(f)(s, t) = f(st) \quad (f \in L^\infty(G), s, t \in G).$$

The left/right Haar measures induce weights.  $L^1(G)$  carries the usual convolution product. On  $L^2(\mathbb{G}) = L^2(G)$  we find that

$$(W\xi)(s, t) = \xi(s, s^{-1}t) \quad (\xi \in L^2(G \times G), s, t \in G).$$

## Duality continued: Fourier algebra

We have  $L^\infty(\widehat{G}) = VN(G)$ , and on the translation operators

$$\widehat{\Delta} : \lambda_s \mapsto \lambda_s \otimes \lambda_s.$$

That such a map exists can be shown by using  $\widehat{W} = \sigma(W^*)$  and that  $\widehat{\Delta}(x) = \widehat{W}^*(1 \otimes x)\widehat{W}$  for  $x \in VN(G)$ .

For general  $\mathbb{G} \dots$  We define  $A(\mathbb{G}) = \widehat{\lambda}(L^1(\widehat{\mathbb{G}}))$  with the norm from  $L^1(\widehat{\mathbb{G}})$ , but thought of as a subalgebra of  $C_0(\mathbb{G})$ .

- Notice that this corresponds to our classical view of  $A(G)$  as a (non-closed) subalgebra of  $C_0(G)$  with norm arising as the predual of  $VN(G)$ .

# Centralisers and Multipliers

Can think of a multiplier of  $A(G)$  as a map  $T : A(G) \rightarrow A(G)$  with  $T(\omega_1\omega_2) = T(\omega_1)\omega_2$ , that is, a module homomorphism.

## Definition

A *left centraliser* of  $L^1(\widehat{\mathbb{G}})$  is a right module homomorphism,  $L(\widehat{\omega}_1 \star \widehat{\omega}_2) = L(\widehat{\omega}_1) \star \widehat{\omega}_2$ .

## Definition

A *left multiplier* of  $A(\mathbb{G})$  is  $a \in L^\infty(\mathbb{G})$  with  $a\widehat{\lambda}(\widehat{\omega}) \in \widehat{\lambda}(L^1(\widehat{\mathbb{G}})) = A(\mathbb{G})$  for each  $\widehat{\omega} \in L^1(\widehat{\mathbb{G}})$ .

As  $\widehat{\lambda}$  is injective, a left multiplier  $a$  induces a (unique) left centraliser  $L$  with  $a\widehat{\lambda}(\widehat{\omega}) = \widehat{\lambda}(L(\widehat{\omega}))$ .

We say that  $L$  (and thus  $a$ ) is *completely bounded* if the adjoint  $L^* : L^\infty(\widehat{\mathbb{G}}) \rightarrow L^\infty(\widehat{\mathbb{G}})$  is completely bounded.

## Centralisers are multipliers

### Theorem (Junge–Neufang–Ruan; D.)

*For any cb left centraliser  $L$  there exists  $a \in M(C_0(\mathbb{G})) \subseteq L^\infty(\mathbb{G})$  an associated multiplier.*

We write  $M_{cb}(A(\mathbb{G}))$  for the collection of all multipliers, equipped with the norm (operator space structure) arising as centralisers, that is, maps on  $L^1(\widehat{\mathbb{G}})$ .

Following the classical situation,  $M_{cb}(A(\mathbb{G}))$  is a dual space: let  $Q_{cb}(A(\mathbb{G}))$  be the closure of the image of  $L^1(\mathbb{G})$  in  $M_{cb}(A(\mathbb{G}))^*$  where  $a \in M_{cb}(A(\mathbb{G}))$  acts on  $\omega \in L^1(\mathbb{G})$  by regarding  $a \in L^\infty(\mathbb{G})$ .

### Definition (D.-Krajczok–Voigt)

$\mathbb{G}$  has the AP if there is a net in  $A(\mathbb{G})$  which converges to 1 weak\* in  $M_{cb}(A(\mathbb{G}))$ .

(We used “left”; there is a “right” analogue; this gives the same notion.)

## Other notions of convergence

Each  $a \in M_{cb}(A(\mathbb{G}))$  is associated to a centraliser  $L : L^1(\widehat{\mathbb{G}}) \rightarrow L^1(\widehat{\mathbb{G}})$  and hence to a map  $L^* = \Theta(a) \in \mathcal{CB}(L^\infty(\widehat{\mathbb{G}}))$ .

### Definition (Crann; Kraus–Ruan)

$\mathbb{G}$  has the (strong) AP when there is a net  $(a_i)$  in  $A(\mathbb{G})$  such  $(\Theta(a_i) \otimes \text{id})(x) \rightarrow x$  weak\* for each  $x \in L^\infty(\widehat{\mathbb{G}}) \bar{\otimes} \mathcal{B}(\ell^2)$  (that is, *stable point-weak\** convergence to id).

### Proposition (DKV)

*AP and strong AP are equivalent.*

### Proof.

Only (AP)  $\implies$  (strong AP) needs a proof. Follows from a careful study of  $Q_{cb}(A(\mathbb{G}))$  and adapting some classical work of Kraus–Haagerup: as sometimes happens you end up proving a little bit more in the abstract setting of LCQGs.  $\square$

# From AP to Operator Algebra APs

## Proposition (Crann; DKV)

*Let  $\mathbb{G}$  have the AP. Then  $L^\infty(\widehat{\mathbb{G}})$  has the  $w^*$ OAP. That is, there is a net of  $w^*$ -cts finite-rank maps  $(\varphi_i)$  on  $L^\infty(\widehat{\mathbb{G}})$  such that  $\varphi_i \otimes \text{id} \rightarrow \text{id}$  pointwise weak\* on  $L^\infty(\widehat{\mathbb{G}}) \bar{\otimes} \mathcal{B}(\ell^2)$ .*

## Proof.

We can assume we have strong AP, so there is a net  $(a_i)$  in  $A(\mathbb{G})$  with  $\Theta(a_i) \rightarrow \text{id}$  in the stable point-weak\* topology on  $L^\infty(\widehat{\mathbb{G}})$ . For each  $i$  there is  $\widehat{\omega}_i \in L^1(\widehat{\mathbb{G}})$  with  $a_i = \widehat{\lambda}(\widehat{\omega}_i)$ .  $\square$

## From AP to Operator Algebra APs (cont.)

### Proof.

Consider  $\widehat{V} \in L^\infty(\mathbb{G})' \bar{\otimes} L^\infty(\widehat{\mathbb{G}})$  the right multiplicative unitary for  $\widehat{\mathbb{G}}$ . Extend each  $\widehat{\omega}_i$  to  $\tilde{\omega}_i \in \mathcal{B}(L^2(\mathbb{G}))_*$ , and define

$$\Psi_i : \mathcal{B}(L^2(\mathbb{G})) \ni x \mapsto (\tilde{\omega}_i \otimes \text{id})(\widehat{V}(x \otimes 1)\widehat{V}^*) \in L^\infty(\widehat{\mathbb{G}}).$$

This map restricts to  $\Theta(a_i)$  on  $L^\infty(\widehat{\mathbb{G}})$ .

These maps nicely approximate the identity, but are not finite-rank. However,  $\mathcal{B}(L^2(\mathbb{G}))$  has the  $w^*$ CPAP, and these provide suitable finite-rank maps, and then we compose to obtain the maps we need. □

## Discrete case

### Proposition (Kraus–Ruan)

*For discrete  $\mathbb{G}$ , consider the following:*

- 1  $\mathbb{G}$  has AP;
- 2  $C(\widehat{\mathbb{G}})$  has the OAP;
- 3  $L^\infty(\widehat{\mathbb{G}})$  has the  $w^*$  OAP

*Then (1) $\Rightarrow$ (2) and (1) $\Rightarrow$ (3) and when  $\mathbb{G}$  is unimodular, all are equivalent.*

## Relative $w^*$ OAP

$M$  has  $w^*$  OAP:  $\varphi_i \rightarrow \text{id}$  stable point- $w^*$ .

Let  $\varphi$  be a weight on  $M$  with GNS space  $L^2(\varphi)$ , definition ideal

$$\mathfrak{n}_\varphi = \{x \in M : \varphi(x^*x) < \infty\},$$

and GNS map  $\Lambda : \mathfrak{n}_\varphi \rightarrow L^2(\varphi)$ . Say that  $\varphi_i$  has an  $L^2$ -implementation when  $\varphi_i(\mathfrak{n}_\varphi) \subseteq \mathfrak{n}_\varphi$ , and there is  $T_i \in \mathcal{B}(L^2(\varphi))$  with  $T_i\Lambda(x) = \Lambda(\varphi_i(x))$  for  $x \in \mathfrak{n}_\varphi$ .

### Definition

Let  $N \subseteq \mathcal{B}(L^2(\varphi))$  be a von Neumann algebra.  $M$  has the  $w^*$  OAP relative to  $N$  when each  $T_i \in N$ .

# Relative $w^*$ OAP and AP

## Theorem (DKV)

For a discrete quantum group  $\mathbb{G}$  the following are equivalent:

- 1  $\mathbb{G}$  has AP;
- 2  $L^\infty(\widehat{\mathbb{G}})$  has  $w^*$  OAP relative to  $\ell^\infty(\mathbb{G})$ ;
- 3  $L^\infty(\widehat{\mathbb{G}})$  has  $w^*$  OAP relative to  $\ell^\infty(\mathbb{G})'$ ;

# Permanence properties

## Theorem (DKV)

*Let  $\mathbb{G}$  have the AP, and let  $\mathbb{H}$  be a closed quantum subgroup of  $\mathbb{G}$ . Then  $\mathbb{H}$  has the AP.*

## Proof.

Almost by definition,  $\mathbb{H} \leq \mathbb{G}$  means that there is a quotient map  $A(\mathbb{G}) \rightarrow A(\mathbb{H})$  (classically this is the Herz Restriction Theorem).  $\square$

## Free products

### Theorem (DKV)

*Let  $\mathbb{G}_1, \mathbb{G}_2$  be discrete quantum groups with the AP. Then  $\mathbb{G}_1 \star \mathbb{G}_2$  has the AP.*

Is there a reference in the classical case?

### Proof.

With  $\mathbb{G} = \mathbb{G}_1 \star \mathbb{G}_2$ , by definition,  $C(\widehat{\mathbb{G}}) = C(\widehat{\mathbb{G}}_1) \star C(\widehat{\mathbb{G}}_2)$ . We use operator algebraic methods to deal with this  $C^*$ -algebraic free product, especially results of [Ricard–Xu]. Then check that these ideas arise (or can be made to arise) from operations on cb-multipliers which are weak\*-continuous. □

## Double crossed product

Let  $\mathbb{G}_1, \mathbb{G}_2$  be locally compact quantum groups. Following [Baaĳ–Vaes], a *matching* is an injective normal  $*$ -homomorphism (which is automatically a  $*$ -isomorphism)

$m : L^\infty(\mathbb{G}_1) \bar{\otimes} L^\infty(\mathbb{G}_2) \rightarrow L^\infty(\mathbb{G}_1) \bar{\otimes} L^\infty(\mathbb{G}_2)$  with

$$(\Delta_1 \otimes \text{id})m = m_{23} m_{13} (\Delta_1 \otimes \text{id}), \quad (\text{id} \otimes \Delta_2)m = m_{13} m_{12} (\text{id} \otimes \Delta_2).$$

From this, we can construct the *double crossed product*  $\mathbb{G}_m$  with

$$L^\infty(\mathbb{G}_m) = L^\infty(\mathbb{G}_1) \bar{\otimes} L^\infty(\mathbb{G}_2), \quad \Delta_m = (\text{id} \otimes \sigma m \otimes \text{id})(\Delta_1^{\text{op}} \otimes \Delta_2).$$

(Notice that the product is a very special case of this.)

## Quantum double: results

### Proposition (DKV)

*If  $\mathbb{G}_m$  has the AP then so do  $\mathbb{G}_1$  and  $\mathbb{G}_2$ .*

### Proof.

$\mathbb{G}_1^{\text{op}}$  and  $\mathbb{G}_2$  are closed quantum subgroups of  $\mathbb{G}_m$ . □

### Theorem (DKV)

*If  $\widehat{\mathbb{G}}_1$  and  $\widehat{\mathbb{G}}_2$  have the AP, then so does  $\widehat{\mathbb{G}}_m$ .*

### Proof.

The idea is to translate the approximating nets from  $A(\widehat{\mathbb{G}}_1)$  and  $A(\widehat{\mathbb{G}}_2)$  to  $A(\widehat{\mathbb{G}}_m)$ . At a key point, this doesn't seem to quite work, but the issue can be side-stepped by using a construction of [Junge–Neufang–Ruan] to extend a centraliser  $\Theta(a) \in \mathcal{CB}(L^\infty(\widehat{\mathbb{G}}))$  to all of  $\mathcal{B}(L^2(\mathbb{G}))$ . □

# Products

## Corollary

*For locally compact quantum groups  $\mathbb{G}_1, \mathbb{G}_2$  the following are equivalent:*

- 1  $\mathbb{G}_1, \mathbb{G}_2$  both have AP;
- 2  $\mathbb{G}_1 \times \mathbb{G}_2$  has AP.

## The (early) end

We would like to know more about when  $\mathbb{G}_m$  has (or does not have) the AP.

Thanks for your attention!