

# Multipliers of locally compact quantum groups and Hilbert $C^*$ -modules

## 3. Quantum groups

Matthew Daws

Leeds

June 2010

## Locally compact quantum groups

Such a thing is a von Neumann algebra  $L^\infty(\mathbb{G})$  together with a unital, injective, normal  $*$ -homomorphism  $\Delta : L^\infty(\mathbb{G}) \rightarrow L^\infty(\mathbb{G}) \overline{\otimes} L^\infty(\mathbb{G})$  such that  $(\Delta \otimes \iota)\Delta = (\iota \otimes \Delta)\Delta$ .

The pre-adjoint  $\Delta_*$  induces an associative product on  $L^1(\mathbb{G}) = L^\infty(\mathbb{G})_*$ . Then  $L^1(\mathbb{G})$  becomes a (completely contractive) Banach algebra.

We assume the existence of left and right normal, faithful, semifinite weights. Using the left weight, we construct a Hilbert space  $L^2(\mathbb{G})$  upon which  $L^\infty(\mathbb{G})$  acts in standard position. There is a unitary map  $W$  (whose existence proof needs the right weight!) such that

$$\Delta(x) = W^*(1 \otimes x)W \quad (x \in L^\infty(\mathbb{G})).$$

If we define  $C_0(\mathbb{G})$  to be the norm closure of

$$\{(\iota \otimes \omega)W : \omega \in B(L^2(\mathbb{G}))_*\}$$

then  $C_0(\mathbb{G})$  is a  $C^*$ -algebra with  $C_0(\mathbb{G})'' = L^\infty(\mathbb{G})$ , and such that  $\Delta$  restricts to a map  $C_0(\mathbb{G}) \rightarrow M(C_0(\mathbb{G}) \otimes C_0(\mathbb{G}))$ . Furthermore, the weights restrict to “nice” weights on  $C_0(\mathbb{G})$ . We think of  $(C_0(\mathbb{G}), \Delta)$  as being the  $C^*$ -algebraic counterpart to  $L^\infty(\mathbb{G})$ .

## Duality

If we let  $C_0(\hat{\mathbb{G}})$  be the norm closure of

$$\{(\omega \otimes \iota)W : \omega \in B(L^2(\mathbb{G}))_*\}$$

then this is a  $C^*$ -algebra; let  $L^\infty(\hat{\mathbb{G}}) = C_0(\hat{\mathbb{G}})''$ . There exists a coassociative  $\hat{\Delta} : L^\infty(\hat{\mathbb{G}}) \rightarrow L^\infty(\hat{\mathbb{G}}) \overline{\otimes} L^\infty(\hat{\mathbb{G}})$ . Also  $L^\infty(\hat{\mathbb{G}})$  admits left and right invariant weights, and so becomes a locally compact quantum group in its own right.

We have that  $\hat{\hat{\mathbb{G}}} = \mathbb{G}$  canonically.

In fact, the map

$$\lambda : L^1(\mathbb{G}) \rightarrow C_0(\hat{\mathbb{G}}); \quad \omega \mapsto (\omega \otimes \iota)W,$$

makes sense, and is a (completely) contractive homomorphism.

We have that:

$$W \in L^\infty(\mathbb{G}) \overline{\otimes} L^\infty(\hat{\mathbb{G}})$$

$$W \in M(C_0(\mathbb{G}) \otimes B_0(L^2(\mathbb{G})))$$

$$W \in M(B_0(L^2(\mathbb{G})) \otimes C_0(\hat{\mathbb{G}}))$$

$$W \in M(C_0(\mathbb{G}) \otimes C_0(\hat{\mathbb{G}})).$$

# Multipliers: from abstract to concrete

## Theorem

Let  $(L, R) \in M_{cb}(L^1(\hat{\mathbb{G}}))$ . There exists  $a \in M(C_0(\mathbb{G}))$  such that

$$\hat{\lambda}(L(\hat{\omega})) = a\hat{\lambda}(\hat{\omega}), \quad \hat{\lambda}(R(\hat{\omega})) = \hat{\lambda}(\hat{\omega})a \quad (\hat{\omega} \in L^1(\hat{\mathbb{G}})).$$

## Proof.

Kraus and Ruan showed this for Kac algebras; not so hard to adapt the ideas to locally compact quantum groups.

Idea is to firstly define  $a$  as a (possibly unbounded) densely defined operator:

$$a\hat{\lambda}(\hat{\omega})\xi = \hat{\lambda}(L(\hat{\omega}))\xi \quad (\xi \in L^2(\mathbb{G}), \hat{\omega} \in L^1(\hat{\mathbb{G}})).$$

That this is well-defined needs the existence of  $R$  (and doesn't use any complete boundedness).

You then show that  $(R^* \otimes \iota)(\hat{W}) = \hat{W}(1 \otimes a)$  on some dense subspace of  $L^2(\mathbb{G}) \otimes L^2(\mathbb{G})$ . As  $\hat{W}$  is unitary, it follows that  $a$  must be bounded. □

# One-sided Multipliers: from abstract to concrete

A corollary of the Junge, Neufang, Ruan representation result is:

## Theorem (JNR)

Let  $R \in M_{cb}^r(L^1(\hat{\mathbb{G}}))$ . Then there exists  $a \in L^\infty(\mathbb{G})$  such that

$$\hat{\lambda}(R(\hat{\omega})) = \hat{\lambda}(\hat{\omega})a \quad (\hat{\omega} \in L^1(\hat{\mathbb{G}})).$$

Here  $M_{cb}^r(L^1(\hat{\mathbb{G}}))$  is the space of completely bounded right multipliers. That is, bounded maps  $R : L^1(\hat{\mathbb{G}}) \rightarrow L^1(\hat{\mathbb{G}})$  such that  $R(ab) = aR(b)$  and with  $R^* : L^\infty(\hat{\mathbb{G}}) \rightarrow L^\infty(\hat{\mathbb{G}})$  being completely bounded.

# Towards a quantum version of Gilbert's Theorem

## Theorem

Let  $f \in C^b(G)$ . The following are equivalent:

- 1  $f \in M_{cb}A(G)$ ;
- 2 there exists a Hilbert space  $K$ , and bounded continuous maps  $\alpha, \beta : G \rightarrow K$ , such that  $f(st^{-1}) = (\alpha(s)|\beta(t))$  for  $s, t \in G$ .

Remember that when  $A = C_0(G)$ ,

$$C_0(G, K) = A \otimes K, \quad C^b(G, K) = \mathcal{L}(A, A \otimes K).$$

In the previous talk, given  $\alpha \in \mathcal{L}(A, A \otimes K)$ , we constructed  $\Delta * \alpha$  (somehow analogous to “applying  $\Delta$  pointwise”) in  $\mathcal{L}(A \otimes A, A \otimes A \otimes K)$ . Then we get:

## Theorem

Let  $f \in M(A)$ . The following are equivalent:

- 1  $f \in M_{cb}A(G)$ ;
- 2 there exists a Hilbert space  $K$  and  $\alpha, \beta \in \mathcal{L}(A, A \otimes K)$  with  $(1 \otimes \beta)^*(\Delta * \alpha) = f \otimes 1$  in  $\mathcal{L}(A \otimes A) = M(A \otimes A)$ .

# On the left

## Theorem

Let  $a \in M(C_0(\mathbb{G}))$ , and let  $L \in \mathcal{CB}(L^1(\hat{\mathbb{G}}))$  be defined by  $L^*(\cdot) = \tilde{\beta}^*(\cdot \otimes 1)\tilde{\alpha}$ , for some  $\alpha, \beta \in \mathcal{L}(C_0(\mathbb{G}), C_0(\mathbb{G}) \otimes K)$ . Then following are equivalent:

- 1  $(1 \otimes \beta)^*(\Delta * \alpha) = a \otimes 1$ ;
- 2  $L$  is a left multiplier, represented by  $a$  in the sense that  $\hat{\lambda}(L(\hat{\omega})) = a\hat{\lambda}(\hat{\omega})$ .

## Proof.

As  $\Delta(\cdot) = W^*(1 \otimes \cdot)W$ , we can recast the first condition as

$$(1 \otimes \tilde{\beta}^*)W_{12}^*(1 \otimes \tilde{\alpha})W = a \otimes 1.$$

Then use that  $\hat{W} = \sigma W^* \sigma$  to show that

$$(L \otimes \iota)(\hat{W}) = (1 \otimes a)\hat{W},$$

which is equivalent to the second condition. (And then reverse the argument).  $\square$

# On the left, harder

## Theorem

Let  $L \in M_{cb}^l(L^1(\hat{G}))$  be represented by  $a \in L^\infty(G)$ . (Always true by [JNR].) Then there exist  $\alpha, \beta \in \mathcal{L}(C_0(G), C_0(G) \otimes K)$  with  $(1 \otimes \beta)^*(\Delta * \alpha) = a \otimes 1$ . So  $a \in M(C_0(G))$ , and  $L^*(x) = \tilde{\beta}^*(x \otimes 1)\tilde{\alpha}$ .

## Proof.

Basic idea is as in Jolissaint's proof of Gilbert's Theorem. We replace a unitary representation of  $G$  by a unitary corepresentation of  $C_0(G)$ . The complication comes that we need to use "universal quantum groups", see the paper of Kustermans. (For example, for non-amenable  $G$ , we really need to work with  $C^*(G)$  and not  $C_r^*(G)$ ). □

## On the right

There exists an antilinear isometry  $J : L^2(\mathbb{G}) \rightarrow L^2(\mathbb{G})$  connected to the Tomita-Takesaki theory of  $L^\infty(\mathbb{G})$ . Remarkably, this gives a map

$$\hat{\kappa} : L^\infty(\hat{\mathbb{G}}) \rightarrow L^\infty(\hat{\mathbb{G}}); \quad x \mapsto Jx^*J$$

which restricts to  $C_0(\hat{\mathbb{G}})$ , and satisfies  $\hat{\Delta}\hat{\kappa} = \sigma(\hat{\kappa} \otimes \hat{\kappa})\hat{\Delta}$ . So the pre-adjoint  $\hat{\kappa}_* : L^1(\hat{\mathbb{G}}) \rightarrow L^1(\hat{\mathbb{G}})$  is an anti-homomorphism. Furthermore,

$$\hat{\lambda} \circ \hat{\kappa}_* = \kappa \circ \hat{\lambda}.$$

### Theorem (Really, a lemma!)

Let  $L$  be a map on  $L^1(\hat{\mathbb{G}})$ , and let  $R = \hat{\kappa}_*L\hat{\kappa}_*$ . Then the following are equivalent:

- 1  $L$  is a completely bounded left multiplier represented by  $a \in L^\infty(\mathbb{G})$  (or in  $M(C_0(\mathbb{G}))$ );
- 2  $R$  is a completely bounded right multiplier represented by  $\kappa(a) \in L^\infty(\mathbb{G})$  (or in  $M(C_0(\mathbb{G}))$ );

So, end of story for right multipliers?

## Using the opposite algebra

Let  $\hat{\mathbb{G}}^{\text{op}}$  be the locally compact quantum group which has  $L^\infty(\hat{\mathbb{G}}^{\text{op}}) = L^\infty(\hat{\mathbb{G}})$  but with  $\hat{\Delta}^{\text{op}} = \sigma\hat{\Delta}$ . We swap the left and right invariant weights.

We have that  $L^1(\hat{\mathbb{G}}^{\text{op}})$  is just the *opposite* Banach algebra to  $L^1(\hat{\mathbb{G}})$ . So left and right multipliers get swapped.

But what is the dual of  $\hat{\mathbb{G}}^{\text{op}}$ ? It is  $\mathbb{G}'$ , where  $L^\infty(\mathbb{G}') = L^\infty(\mathbb{G})'$ , the commutant. Tomita theory tells us that  $L^\infty(\mathbb{G}') = JL^\infty(\mathbb{G})J$ . We can similarly build a coproduct and weights. The multiplicative unitary is

$$W' = (J \otimes J)W(J \otimes J),$$

from which it follows that also  $C_0(\mathbb{G}') = JC_0(\mathbb{G})J$ .

There is a bijection between  $\alpha' \in \mathcal{L}(C_0(\mathbb{G}'), C_0(\mathbb{G}') \otimes K)$  and  $\alpha \in \mathcal{L}(C_0(\mathbb{G}), C_0(\mathbb{G}) \otimes K)$  (we need to pick an “involution”  $J_K$  on  $K$ , so this isn't totally canonical).

Then  $(1 \otimes \beta')^*(\Delta' * \alpha') = JaJ \otimes 1$  if and only if  $(1 \otimes \beta)^*(\Delta * \alpha) = a \otimes 1$ .

## Swapping things about

Pick  $R \in M_{cb}^r(L^1(\hat{\mathbb{G}}))$ , and consider  $R^{\text{op}} \in M_{cb}^l(L^1(\hat{\mathbb{G}}^{\text{op}}))$ . Let  $a$  and  $JbJ$  “represent”  $R$  and  $R^{\text{op}}$ :

$$\hat{\lambda}(R(\hat{\omega})) = \hat{\lambda}(\hat{\omega})a, \quad \hat{\lambda}^{\text{op}}(R^{\text{op}}(\hat{\omega})) = JbJ\hat{\lambda}^{\text{op}}(\hat{\omega}).$$

We can find  $\alpha', \beta'$  associated to  $R^{\text{op}}$ ; this leads to  $\alpha, \beta$  with

$$b \otimes 1 = (1 \otimes \beta)^*(\Delta * \alpha).$$

It turns out that

$$\kappa(a) \otimes 1 = (1 \otimes \alpha)^*(\Delta * \beta).$$

We call a pair  $(\alpha, \beta)$  “invariant” if  $(1 \otimes \beta)^*(\Delta * \alpha) \in M(C_0(\mathbb{G})) \otimes 1$ .

So we’ve naturally found that  $(\alpha, \beta)$  is invariant (say for  $b$ ) if and only if  $(\beta, \alpha)$  is invariant (say for  $\kappa(a)$ ). What’s the relationship between  $a$  and  $b$ ?

# The (unbounded) antipode

The *antipode*  $S$  on  $\mathbb{G}$  is an unbounded operator: it plays the role of the inverse in quantum group theory. For Kac algebras, we have that  $S = \kappa$ , but even for compact quantum groups,  $S$  may be unbounded and only densely defined. For example, we have

$$S((\iota \otimes \omega)(W)) = (\iota \otimes \omega)(W^*) \quad (\omega \in B(H)_*).$$

We can “split off” the unbounded part of  $S$ . There is a strong\*-continuous automorphism group  $(\sigma_t)_{t \in \mathbb{R}}$  of  $L^\infty(\mathbb{G})$  (which restricts to a norm continuous automorphism group of  $C_0(\mathbb{G})$ ). You can analytically extend this to unbounded maps; then

$$S = \kappa \tau_{-i/2}.$$

Then, if  $(\alpha, \beta) \rightarrow b$  and  $(\beta, \alpha) \rightarrow \kappa(a)$ , then  $b = \tau_{-i/2}(a^*)$ .

## Taking a coordinate approach

Let  $(e_i)$  be an orthonormal basis for  $K$ . Recall that we can view  $C_0(\mathbb{G}) \otimes K$  as being those families  $(a_i)$  in  $C_0(\mathbb{G})$  with  $\sum_i a_i^* a_i$  converging in norm.

Similarly, we can view  $\mathcal{L}(C_0(\mathbb{G}), C_0(\mathbb{G}) \otimes K)$  as being those families  $(a_i)$  in  $\mathcal{L}(C_0(\mathbb{G})) = M(C_0(\mathbb{G}))$  with  $\sum_i a_i^* a_i$  converging in the *strict* topology.

So let  $\alpha$  be associated to  $(\alpha_i)$ , and similarly for  $\beta$ . Then  $(\alpha, \beta)$  is invariant for  $b$  if and only if

$$\sum_i (1 \otimes \beta_i^*) \Delta(\alpha_i) = b \otimes 1.$$

### Theorem

Let  $\bar{S}$  be the strict extension of  $S$  to  $M(C_0(\mathbb{G}))$ . If  $a \in M(C_0(\mathbb{G}))$  and we have families  $(\alpha_i), (\beta_i)$  with

$$a \otimes 1 = \sum_i \Delta(\alpha_i)(1 \otimes \beta_i),$$

then  $a \in D(\bar{S})$  and

$$\bar{S}(a) \otimes 1 = \sum_i (1 \otimes \alpha_i) \Delta(\beta_i).$$

# Universal quantum groups

For a group  $G$ , we let  $C^*(G)$  be the universal (or full) group  $C^*$ -algebra: this is the completion of  $L^1(G)$  under the largest  $C^*$ -norm.

We can do a similar thing for quantum groups: but firstly we need to turn  $L^1(\mathbb{G})$  into a  $*$ -algebra. The standard way to do this is to use the involution, so we need to restrict to a subalgebra  $L^1_{\#}(\mathbb{G})$  of  $L^1(\mathbb{G})$  where this is bounded. This then leads to  $C_u(\hat{\mathbb{G}})$ . Denote the  $*$ -representation by

$$\lambda_u : L^1_{\#}(\mathbb{G}) \rightarrow C_u(\hat{\mathbb{G}}).$$

The dual of  $C_u(\hat{\mathbb{G}})$  is a dual Banach algebra (follow an argument of [Runde]) which contains  $L^1(\hat{\mathbb{G}})$  as an ideal. So we get an inclusion  $C_u(\hat{\mathbb{G}})^* \rightarrow M_{cb}(L^1(\hat{\mathbb{G}}))$ .

**Aside:** If  $\hat{\mathbb{G}}$  is *co-amenable* ( $L^1(\hat{\mathbb{G}})$  has a bai) then  $C_u(\hat{\mathbb{G}}) = C_0(\hat{\mathbb{G}})$  and  $M_{cb}(L^1(\hat{\mathbb{G}})) = C_0(\hat{\mathbb{G}})^*$ . Is the converse true? (This is Losert's Theorem for  $A(G)$ .)

# Universal quantum groups: in our framework

Pick  $\mu \in C_u(\hat{\mathbb{G}})^*$  and choose a representation  $\theta : C_u(\hat{\mathbb{G}}) \rightarrow B(K)$  and  $\xi, \eta \in K$  with

$$\mu(a) = (\xi | \theta(a) \eta) \quad (a \in C_u(\hat{\mathbb{G}})).$$

[Kustermans]  $\Rightarrow$  there is a unitary  $U \in M(C_0(\mathbb{G}) \otimes B_0(K))$  with

$$\theta(\lambda_u(\omega)) = (\omega \otimes \iota)(U), \quad (\Delta \otimes \iota)(U) = U_{13}U_{23}.$$

Associate  $U$  with  $\mathcal{U} \in \mathcal{L}(C_0(\mathbb{G}) \otimes K)$ , and let

$$\alpha = \mathcal{U}^*(\iota \otimes \xi)^*, \quad \beta = \mathcal{U}^*(\iota \otimes \eta)^*.$$

Then  $(\alpha, \beta)$  induces the left multiplier induced by  $\mu$ . (Indeed, if  $\mu$  were actually in  $L^1(\hat{\mathbb{G}})$  already, we could take  $U = W$ .)

# Double multipliers

## Theorem

Let  $(L, R) \in M_{cb}(L^1(\hat{\mathbb{G}}))$ . There exists a Hilbert space  $K$  with an involution  $J_K$ ,  $\mathcal{T} \in \mathcal{L}(C_0(\mathbb{G}) \otimes K)$  and  $\xi, \eta \in K$  such that:

- with  $\alpha = \mathcal{T}(\iota \otimes \xi)^*$  and  $\beta = \mathcal{T}(\iota \otimes \eta)^*$ , we have that  $(\alpha, \beta)$  induces  $L$ ;
- with  $\alpha = \mathcal{T}(\iota \otimes J_K \eta)^*$  and  $\beta = \mathcal{T}(\iota \otimes J_K \xi)^*$ , we have that  $(\alpha, \beta)$  induces  $\hat{\kappa}_* R \hat{\kappa}_*$  (and thus  $R$ ).

## Proof.

The proof “glues” two Hilbert spaces together, but this isn’t entirely trivial: you definitely need that  $(L, R)$  is a double multiplier (and not just unconnected left and right multipliers).. □

If  $(L, R)$  is induced by  $C_u(\hat{\mathbb{G}})^*$ , then we can take  $\mathcal{T}$  to be unitary. Is the converse true? (Probably equivalent to my earlier question!)