

# Multipliers of locally compact quantum groups and Hilbert $C^*$ -modules

## 2. Multipliers and Hilbert $C^*$ -modules

Matthew Daws

Leeds

June 2010

### Multipliers of $L^1(G)$

We identify  $M(G)$  with the dual of  $C_0(G)$ , and then our coproduct  $\Delta$  induces a product on  $M(G)$ :

$$\langle \mu * \lambda, f \rangle = \int_{G \times G} f(st) d\mu(s) d\lambda(t) \quad (f \in C_0(G), \mu, \lambda \in M(G)).$$

#### Theorem

For any locally compact group  $G$ , we have an isometric isomorphism between  $M(L^1(G))$  and  $M(G)$ .

#### Proof.

We embed  $L^1(G)$  into  $C_0(G)^* = M(G)$  by integration. Then  $L^1(G)$  is an ideal in  $M(G)$ , so we get a contraction  $M(G) \rightarrow M(L^1(G))$ . Conversely, given  $(L, R) \in M(L^1(G))$ , let  $(e_\alpha)$  be a cai for  $L^1(G)$ , and define  $\mu \in M(G)$  to be the weak\*-limit of  $L(e_\alpha)$ . □

## Multipliers of Banach algebras

For a Banach algebra  $A$ , we always assume that  $A$  is *faithful*: if  $a \in A$  and  $bac = 0$  for all  $b, c \in A$ , then  $a = 0$ .

Recall that the *multiplier algebra* is  $M(A)$ , consisting of pairs of maps  $(L, R)$  with  $aL(b) = R(a)b$  for  $a, b \in A$ . If  $A$  is Arens regular and has a bounded approximate identity,

$$M(A) = \{x \in A^{**} : xa, ax \in A \ (a \in A)\}.$$

Homework: Let  $\theta : A \rightarrow B(A)$  be the left-regular representation,  $\theta(a) : b \mapsto ab$ . Show that we can identify  $M(A)$  with

$$\{T \in B(A) : T\theta(a), \theta(a)T \in \theta(A) \ (a \in A)\}.$$

For a locally compact group  $G$ , the algebra  $L^1(G)$  always has a contractive approximate identity (so is faithful). The algebra  $A(G)$  has a bounded approximate identity only when  $G$  is amenable, but is always faithful.

### Multipliers of $A(G)$

#### Theorem

Let  $A \subseteq C_0(G)$  be a sub-algebra such that:

- $A$  is a Banach algebra for some norm such that the inclusion  $A \rightarrow C_0(G)$  is continuous;
- for each  $s \in G$  there exists  $a \in A$  with  $a(s) = 1$  and  $\|a\| \leq 2$ ;
- for each  $s \in G$  there is an open set  $U$  containing  $s$ , and  $a \in A$  with  $a|_U \equiv 1$ .

Then we can identify  $M(A)$  with  $\{f \in C^b(G) : fa \in A \ (a \in A)\}$ .

#### Proof.

As  $A$  is commutative,  $M(A) = \{T : A \rightarrow A : T(ab) = T(a)b \ (a, b \in A)\}$ . The rest is homework. □

Observe that  $A(G)$  satisfies all these conditions. Observe that  $M(C_{00}(G)) = C(G)$ , the algebra of all continuous functions.

## Completely bounded multipliers of $A(G)$

We see that

$$\begin{aligned} M(A(G)) &= \{T \in B(A(G)) : T(ab) = T(a)b \ (a, b \in A(G))\} \\ &= \{f \in C^b(G) : fa \in A(G) \ (a \in A(G))\}. \end{aligned}$$

Given  $T \in M(A(G))$ , we see that  $T^* \in B(VN(G))$ . Hence

$$I \otimes T^* : M_n \otimes VN(G) = M_n(VN(G)) \rightarrow M_n(VN(G)),$$

where  $M_n(VN(G))$  is a von Neumann algebra acting on  $\ell_n^2 \otimes L^2(G)$ .

We say that  $T$  is *completely bounded* if  $I \otimes T^*$  is bounded, uniformly in  $n \in \mathbb{N}$ . Write  $M_{cb}A(G)$  for the algebra of such multipliers.

So, formally,  $M_{cb}A(G)$  is a subalgebra of  $C^b(G)$ . Can we find a characterisation which doesn't involve maps on  $VN(G)$ ?



## Completely bounded maps

Remember that a map  $T$  from a  $C^*$ -algebra  $A$  to  $B(H)$  is completely bounded if and only if we can find a  $*$ -representation  $\pi : A \rightarrow B(K)$  and bounded maps  $P, Q : H \rightarrow K$  with

$$T(x) = Q^* \pi(x) P \quad (x \in A).$$

So, suppose we have  $\alpha, \beta : G \rightarrow K$  bounded and continuous with

$$f(st^{-1}) = (\alpha(s)|\beta(t)) \quad (s, t \in G).$$

Then define  $\tilde{\alpha} : L^2(G) \rightarrow L^2(G, K) = L^2(G) \otimes K$  by  $\tilde{\alpha}(\xi) = (\xi(r)\alpha(r))_{r \in G}$ .

Notice then that

$$(\lambda(s) \otimes 1)\tilde{\alpha}(\xi) = (\xi(s^{-1}r)\alpha(s^{-1}r))_{r \in G} \quad (s \in G).$$

If we form  $\tilde{\beta}$  similarly, then we can define a completely bounded map  $T : VN(G) \rightarrow B(L^2(G))$  by

$$T(x) = \tilde{\beta}^*(x \otimes 1)\tilde{\alpha} \quad (x \in VN(G)).$$

Notice that  $T$  is clearly normal.



## Gilbert's Theorem

### Theorem

Let  $f \in C^b(G)$ . The following are equivalent:

- 1  $f \in M_{cb}A(G)$ ;
- 2 there exists a Hilbert space  $K$ , and bounded continuous maps  $\alpha, \beta : G \rightarrow K$ , such that  $f(st^{-1}) = (\alpha(s)|\beta(t))$  for  $s, t \in G$ .

### Proof.

See Jolissaint, 1992. (History: Jolissaint points us to Cowling and Haagerup, Inventiones, 1989. They point us to Bozejko and Fendler, 1984, who attribute the result to Gilbert, unpublished, late 70s).  $\square$



## Easy direction continued

Then for  $\xi, \eta \in L^2(G)$  and  $s \in G$ ,

$$\begin{aligned} (T(\lambda(s^{-1}))\xi|\eta) &= (\tilde{\beta}^*(\lambda(s^{-1}) \otimes 1)\tilde{\alpha}\xi|\eta) \\ &= \int_G (\xi(sr)\alpha(sr)|\eta(r)\beta(r)) \, dr \\ &= \int_G \xi(sr)\overline{\eta(r)}(\alpha(sr)|\beta(r)) \, dr \\ &= f(srr^{-1}) \int_G \xi(sr)\overline{\eta(r)} \, dr = f(s)(\lambda(s^{-1})\xi|\eta). \end{aligned}$$

Thus  $T(\lambda(s^{-1})) = f(s)\lambda(s^{-1})$ . As  $T$  is normal, it follows that  $T$  maps into  $VN(G)$ .

Remember that  $A(G) \rightarrow C_0(G)$  is the map  $\omega_{\xi, \eta} \mapsto ((\lambda(s^{-1})\xi|\eta))_{s \in G}$ . It follows that  $T$  is the adjoint of the multiplier induced by  $f$ . Thus  $f$  is completely bounded.



## The converse

Now suppose that  $f$  is completely bounded, say inducing  $T : VN(G) \rightarrow VN(G)$  which has the form

$$T(x) = Q^* \pi(x) P \quad (x \in VN(G)),$$

where  $\pi : VN(G) \rightarrow B(K)$  is a  $*$ -representation. As  $T$  is normal, we may suppose that  $\pi$  is too. Notice that we may assume that  $\pi(1) = 1$ . Then the map  $\sigma : G \rightarrow B(K); s \mapsto \pi(\lambda(s))$  is a continuous unitary representation. Now notice that for  $s \in G$  and  $\omega \in A(G)$ ,

$$f(s) \langle \lambda(s^{-1}), \omega \rangle = \langle \lambda(s^{-1}), f\omega \rangle = \langle T(\lambda(s^{-1})), \omega \rangle = \langle Q^* \pi(\lambda(s^{-1})) P, \omega \rangle,$$

and so  $Q^* \sigma(s) P = f(s^{-1}) \lambda(s)$ . Pick  $\xi_0 \in L^2(G)$  a unit vector, and define

$$\alpha(s) = \sigma(s^{-1}) P \lambda(s) \xi_0, \quad \beta(s) = \sigma(s^{-1}) Q \lambda(s) \xi_0 \quad (s \in G).$$

Thus, for  $s, t \in G$ ,

$$\begin{aligned} \langle \alpha(s) | \beta(t) \rangle &= \langle Q^* \sigma(t^{-1})^* \sigma(s^{-1}) P \lambda(s) \xi_0 | \lambda(t) \xi_0 \rangle \\ &= \langle f(st^{-1}) (\lambda(ts^{-1}) \lambda(s) \xi_0) | \lambda(t) \xi_0 \rangle = \langle f(st^{-1}) \xi_0 | \xi_0 \rangle. \end{aligned}$$



## Examples

- If  $A = \mathbb{C}$ , then we just recover the notion of a Hilbert space.
- Given  $A$ , we can turn  $A$  into a Hilbert  $C^*$ -module over itself by defining  $\langle a | b \rangle = a^* b$ .
- Let  $K$  be a Hilbert space, and form the algebraic tensor product  $A \odot K$ . This becomes a pre-Hilbert  $C^*$ -module for

$$\langle a \otimes \xi | b \otimes \eta \rangle = a^* b \langle \xi | \eta \rangle,$$

with the module action  $(a \otimes \xi) \cdot b = ab \otimes \xi$ .

To show that this is positive definite, notice that we can write any tensor in  $A \odot K$  as  $\sum a_k \otimes \xi_k$  with the  $(\xi_k)$  being orthonormal. Then the inner-product is  $\sum_k a_k^* a_k \geq 0$ .

Let  $A \otimes K$  be the completion of  $A \odot K$ .

Indeed, if  $(e_i)$  is an orthonormal basis for  $K$ , then  $A \otimes K$  consists of those families  $(a_i)$  in  $A$  such that  $\sum_i a_i^* a_i$  converges in  $A$  (notice that this is weaker than  $\sum_i \|a_i\|^2 < \infty$ ).



## Hilbert $C^*$ -modules

We have seen that the space  $C^b(G, K)$  is very useful. How can we think about this abstractly?

Let  $A$  be a  $C^*$ -algebra. A *pre-Hilbert  $C^*$ -module* over  $A$  is a *right  $A$ -module*  $E$  where  $E$  admits an  $A$ -valued sesquilinear map satisfying:

- 1  $\langle x | \cdot \rangle$  is a *linear* map  $E \rightarrow A$  and  $\langle x | y \rangle^* = \langle y | x \rangle$ ;
- 2  $\langle x | x \rangle \geq 0$  in the  $C^*$ -algebra sense, and  $\langle x | x \rangle = 0 \implies x = 0$ ;
- 3  $\langle x | y \cdot a \rangle = \langle x | y \rangle a$ .

Henceforth all inner-products will be linear on the right.

We can define a norm on  $E$  by  $\|x\| = \|\langle x | x \rangle\|_A^{1/2}$ ; this is a norm, which follows by first showing a Cauchy-Schwarz inequality for  $\langle \cdot | \cdot \rangle$ . When  $E$  is complete, we say that  $E$  is *Hilbert  $C^*$ -module*.

Notice that

$$\|x \cdot a\|^2 = \|\langle x \cdot a | x \cdot a \rangle\| = \|\langle x | x \cdot a \rangle^* a\| = \|a^* \langle x | x \rangle a\| \leq \|\langle x | x \rangle\| \|a^* a\| = \|x\|^2 \|a\|^2.$$



## Morphisms

Let  $E$  and  $F$  be Hilbert  $C^*$ -modules over  $A$ . A map  $T : E \rightarrow F$  is *adjointable* if there exists a map  $T^* : F \rightarrow E$  with

$$\langle T(x) | y \rangle = \langle x | T^*(y) \rangle \quad (x \in E, y \in F).$$

Homework: Show that  $T$  and  $T^*$  are linear, and using the Closed Graph Theorem, that  $T$  and  $T^*$  are bounded. Show furthermore that  $T$  and  $T^*$  are  $A$ -module maps.

Unlike for Hilbert spaces, if  $T$  is bounded and linear (and an  $A$ -module map) then we cannot always find  $T^*$ .

Clearly  $T^{**} = T$ , and it's easy to see that  $\|T^* T\| = \|T\|^2$ . We write  $\mathcal{L}(E, F)$  for the collection of adjointable maps  $E \rightarrow F$ . Then  $\mathcal{L}(E)$  is a  $C^*$ -algebra.

Consider the "finite-rank" map

$$\theta_{x,y} : E \rightarrow F; z \mapsto x \cdot \langle y | z \rangle \quad (z \in E),$$

where  $x \in F$  and  $y \in E$ . This is adjointable, because  $\theta_{x,y}^* = \theta_{y,x}$ . Let  $\mathcal{K}(E)$  be the closure of the linear span of such maps in  $\mathcal{L}(E, F)$ .



## Links with multipliers

### Theorem

Consider  $A$  as a module over itself. Then  $\mathcal{K}(A) \cong A$  and  $\mathcal{L}(A) \cong M(A)$ .

### Proof.

Let  $a, b \in A$ , so that  $\theta_{a,b}(c) = a|b|c = ab^*c$ , and hence  $\theta_{a,b}$  is left multiplication by  $ab^*$ . As  $A$  has an approximate identity, it's not hard to see that  $\mathcal{K}(A) \cong A$ . Given  $T \in \mathcal{L}(A)$ , define  $(L, R) \in M(A)$  by

$$L(a) = T(a), \quad R(a) = T^*(a^*)^* \quad (a \in A).$$

Then  $a^*L(b) = (a|T(b)) = (T^*(a)|b) = T^*(a^*)b = R(a^*)b$ . So we have a map  $\mathcal{L}(A) \rightarrow M(A)$ . This is onto, as given  $(L, R) \in M(A)$ , the map  $L$  is adjointable, with  $L^*(a) = R(a^*)^*$ .  $\square$

For general  $E$ , we have that  $\mathcal{K}(E)$  is an ideal in  $\mathcal{L}(E)$ , and, considering  $\mathcal{K}(E)$  as a  $C^*$ -algebra, we have that  $M(\mathcal{K}(E)) \cong \mathcal{L}(E)$ .



## Links with continuous functions

### Theorem

We can identify  $C_0(G) \otimes K$  with  $C_0(G, K)$ .

### Proof.

Algebraically, we identify  $f \otimes \xi$  with  $(f(s)\xi)_{s \in G}$ . Then

$$\begin{aligned} \left\| \sum_k f_k \otimes \xi_k \right\|^2 &= \left\| \sum_{j,k} f_j^* f_k (\xi_j | \xi_k) \right\|_{C_0(G)} = \sup_{s \in G} \left| \sum_{j,k} \overline{f_j(s)} f_k(s) (\xi_j | \xi_k) \right| \\ &= \sup_{s \in G} \left| \sum_{j,k} (f_j(s)\xi_j | f_k(s)\xi_k) \right| = \sup_{s \in G} \left\| \sum_k f_k(s)\xi_k \right\|^2. \end{aligned}$$

Finally, a partition of unity argument shows that the image of  $C_0(G) \odot K$  is dense in  $C_0(G, K)$ .  $\square$



## For morphisms

### Theorem

We can identify  $\mathcal{L}(C_0(G), C_0(G) \otimes K)$  with  $C^b(G, K)$ .

### Proof.

Given  $F \in C^b(G, K)$ , we define  $T : C_0(G) \rightarrow C_0(G, K)$  by  $T(f) = (f(s)F(s))_{s \in G}$ . This is adjointable, with  $T^*(g) = ((F(s)|g(s)))_{s \in G}$  for  $g \in C_0(G, K)$ .

Conversely, given  $T : C_0(G) \rightarrow C_0(G, K)$  adjointable, notice that

$$T^*(g)(s)f(s) = (T^*(g)|f)(s) = (g|T(f))(s) = (g(s)|T(f)(s)) \quad (s \in G).$$

It follows that if  $f_1(s) = f_2(s) = 1$ , then  $(g(s)|T(f_1)(s)) = (g(s)|T(f_2)(s))$  for all  $g$ ; thus  $T(f_1)(s) = T(f_2)(s)$ . Let this be  $F(s)$ , so  $T(f) = fF$ . Similar arguments show that  $F$  is continuous, and bounded.  $\square$



## More links between multipliers and adjointable maps

### Theorem

We have that  $\mathcal{K}(A \otimes K) \cong A \otimes_{\min} B_0(K)$  as  $C^*$ -algebras, and thus  $\mathcal{L}(A \otimes K) \cong M(A \otimes_{\min} B_0(K))$ .

### Proof.

Algebraically, we define

$$\Theta : \theta_{a \otimes \xi, b \otimes \eta} \mapsto ab^* \otimes \theta_{\xi, \eta},$$

where  $\theta_{\xi, \eta} : K \rightarrow K; \gamma \mapsto \xi(\eta|\gamma)$ , so  $\theta_{\xi, \eta} \in B_0(K)$ . It's easy to see that this defines a homomorphism between a dense subalgebra of  $\mathcal{K}(A \otimes K)$  and a dense subalgebra of  $A \otimes B_0(K)$ .

So it remains to show that  $\Theta$  is an isometry (or at least bounded) and so extends by continuity: this I believe is tricky! (See [Lance]).  $\square$



## Back to groups

Remember that we have  $W \in M(C_0(G) \otimes B_0(H))$  for  $H = L^2(G)$ ,

$$W\xi(s, t) = \xi(s, s^{-1}t) \quad (s, t \in G).$$

Thus  $W$  induces some

$$\mathcal{W} \in \mathcal{L}(C_0(G) \otimes H) = \mathcal{L}(C_0(G, H)).$$

Homework: follow the isomorphisms through to show that

$$\mathcal{W} : f \mapsto (\lambda(s)f(s))_{s \in G} \quad (f \in C_0(G, H)).$$

(Indeed, if  $H$  is any Hilbert space, and  $\sigma$  is a unitary representation of  $G$  on  $H$ , then we can define  $\mathcal{V} \in \mathcal{L}(C_0(G, H))$  in the same way  $f \mapsto (\sigma(s)f(s))_{s \in G}$ . This induces a unitary  $V \in M(C_0(G) \otimes B_0(H))$ : this is the notion of a *corepresentation*.)

## Applications

Let  $\alpha \in \mathcal{L}(A, A \otimes K)$  and  $\mathcal{T} \in \mathcal{L}(A \otimes K)$  be related by  $\alpha = \mathcal{T}(\iota \otimes \xi_0)^*$ . Let  $A \subseteq B(H)$ . As  $\mathcal{L}(A \otimes K) \cong M(A \otimes B_0(K))$ , let  $\mathcal{T}$  be related to  $T \in M(A \otimes B_0(K)) \subseteq B(H \otimes K)$ .

- We can define an operator  $\tilde{\alpha} : H \rightarrow H \otimes K$  by  $\tilde{\alpha}(\xi) = \mathcal{T}(\xi \otimes \xi_0)$ . Then:
  - ▶  $\tilde{\alpha}$  only depends upon  $\alpha$  (and not  $\mathcal{T}$ ).
  - ▶ In fact,  $\tilde{\alpha}^* \tilde{\alpha} = \alpha^* \alpha \in \mathcal{L}(A) \cong M(A) \subseteq B(H)$ .
  - ▶ This is the generalisation of the way we moved from  $C^b(G, K)$  to  $\mathcal{B}(L^2(G), L^2(G, K))$ .
- Given a non-degenerate  $*$ -homomorphism  $\phi : A \rightarrow B$ , we can extend  $\phi \otimes \iota$  to multiplier algebras, and hence define  $S = (\phi \otimes \iota)T \in M(B \otimes B_0(K))$ . Form  $S$  using  $S$ , and let  $\phi * \alpha = S(\iota \otimes \xi_0)^* \in \mathcal{L}(B, B \otimes K)$ . Then:
  - ▶  $\phi * \alpha$  only depends upon  $\alpha$ , not  $\mathcal{T}$ .
  - ▶ indeed, for any  $\xi \in K$ , we have that  $(\iota \otimes \xi)(\phi * \alpha) = \phi((\iota \otimes \xi)\alpha)$  in  $\mathcal{L}(B)$ .

## Slicing

Given  $\xi \in K$ , we can define

$$\iota \otimes \xi : A \otimes K \rightarrow A; \quad a \otimes \eta \mapsto a(\xi|\eta).$$

This is adjointable with

$$(\iota \otimes \xi)^* : A \rightarrow A \otimes K; \quad a \mapsto a \otimes \xi.$$

(This requires a small amount of work: homework!)

Fix a unit vector  $\xi_0 \in K$ . Using these maps, we can view  $\mathcal{L}(A, A \otimes K)$  as a complemented submodule of  $\mathcal{L}(A \otimes K)$ :

$$\begin{aligned} \mathcal{L}(A, A \otimes K) &\rightarrow \mathcal{L}(A \otimes K); & \alpha &\mapsto \alpha(\iota \otimes \xi_0), \\ \mathcal{L}(A \otimes K) &\rightarrow \mathcal{L}(A, A \otimes K); & \mathcal{T} &\mapsto \mathcal{T}(\iota \otimes \xi_0)^*, \end{aligned}$$

which follows, as  $(\iota \otimes \eta)(\iota \otimes \xi)^* = (\eta|\xi) \text{id}_A$ .

## Example

For example, we can apply the second construction to  $\Delta$  to give a map

$$\mathcal{L}(C_0(G), C_0(G) \otimes K) \rightarrow \mathcal{L}(C_0(G \times G), C_0(G \times G) \otimes K); \quad \alpha \mapsto \Delta * \alpha.$$

This is just the map

$$C^b(G, K) \rightarrow C^b(G \times G, K); \quad \alpha \mapsto (\alpha(st))_{(s,t) \in G \times G}.$$

Gilbert's theorem asks that

$$f(st^{-1}) = (\beta(t)|\alpha(s)) \quad (s, t \in G).$$

(Remember: inner-products are linear on the right now!) This is equivalent to

$$f(r) = (\beta(s)|\alpha(rs)) \quad (s, r \in G).$$

In our abstract language, we get

$$(1 \otimes \beta)^*(\Delta * \alpha) = f \otimes 1 \in M(C_0(G) \otimes C_0(G)) = C^b(G \times G).$$