

Multipliers and Abstract Harmonic Analysis

Matthew Daws

Leeds

August 2010

Outline

- 1 Multiplier algebras; Dual Banach algebras
- 2 The Fourier algebra; Extending homomorphisms
- 3 Hopf convolution algebras

Multiplier algebras

Let A be an algebra. A multiplier of A is a pair (L, R) of maps $A \rightarrow A$ such that $aL(b) = R(a)b$ for $a, b \in A$. Let $M(A)$ be the collection of such maps, made into an algebra for the product $(L, R)(L', R') = (LL', R'R)$.

Henceforth assume that A is *faithful*: if $a \in A$ and $bac = 0$ for all $b, c \in A$, then $a = 0$. Then we can show that

$$L(ab) = L(a)b, \quad R(ab) = aR(b) \quad (a, b \in A),$$

and furthermore, the map $A \rightarrow M(A)$,

$$a \mapsto (L_a, R_a), \quad L_a(b) = ab, R_a(b) = ba \quad (a, b \in A),$$

is an injective algebra homomorphism.

Then A becomes an ideal in $M(A)$. If B is an algebra containing A as an ideal, we say that A is *essential* if $x \in B$ is such that $axb = 0$ for $a, b \in A$, then $x = 0$. Then B embeds into $M(A)$. In this sense, $M(A)$ is the “largest” algebra containing A as an essential ideal.

Multiplier algebras

Let A be an algebra. A multiplier of A is a pair (L, R) of maps $A \rightarrow A$ such that $aL(b) = R(a)b$ for $a, b \in A$. Let $M(A)$ be the collection of such maps, made into an algebra for the product $(L, R)(L', R') = (LL', R'R)$.

Henceforth assume that A is *faithful*: if $a \in A$ and $bac = 0$ for all $b, c \in A$, then $a = 0$. Then we can show that

$$L(ab) = L(a)b, \quad R(ab) = aR(b) \quad (a, b \in A),$$

and furthermore, the map $A \rightarrow M(A)$,

$$a \mapsto (L_a, R_a), \quad L_a(b) = ab, R_a(b) = ba \quad (a, b \in A),$$

is an injective algebra homomorphism.

Then A becomes an ideal in $M(A)$. If B is an algebra containing A as an ideal, we say that A is *essential* if $x \in B$ is such that $axb = 0$ for $a, b \in A$, then $x = 0$. Then B embeds into $M(A)$. In this sense, $M(A)$ is the “largest” algebra containing A as an essential ideal.

Multiplier algebras

Let A be an algebra. A multiplier of A is a pair (L, R) of maps $A \rightarrow A$ such that $aL(b) = R(a)b$ for $a, b \in A$. Let $M(A)$ be the collection of such maps, made into an algebra for the product $(L, R)(L', R') = (LL', R'R)$.

Henceforth assume that A is *faithful*: if $a \in A$ and $bac = 0$ for all $b, c \in A$, then $a = 0$. Then we can show that

$$L(ab) = L(a)b, \quad R(ab) = aR(b) \quad (a, b \in A),$$

and furthermore, the map $A \rightarrow M(A)$,

$$a \mapsto (L_a, R_a), \quad L_a(b) = ab, R_a(b) = ba \quad (a, b \in A),$$

is an injective algebra homomorphism.

Then A becomes an ideal in $M(A)$. If B is an algebra containing A as an ideal, we say that A is *essential* if $x \in B$ is such that $axb = 0$ for $a, b \in A$, then $x = 0$. Then B embeds into $M(A)$. In this sense, $M(A)$ is the “largest” algebra containing A as an essential ideal.

Banach algebras; Examples

If A is a Banach algebra, then a little closed graph argument shows that if $(L, R) \in M(A)$, then L and R are bounded. We norm $M(A)$ by regarding it as a subspace of $\mathcal{B}(A) \oplus_{\infty} \mathcal{B}(A)$.

If A is unital, then $A = M(A)$.

If A is a C^* -algebra then so is $M(A)$. For a commutative C^* -algebra $A = C_0(X)$, the multiplier algebra can be identified with $C^b(X)$, which in turn is $C(\beta X)$. So multiplier algebras are Stone-Cech compactifications.

Notice that $M(A)$ is rarely a von Neumann algebra.

Let E be a Banach algebra, and $A = \mathcal{K}(E)$ the compact operators on E . Then $M(A) = \mathcal{B}(E)$.

Notice that $\mathcal{B}(E)$ may or may not be a dual space.

For a locally compact group G , consider the algebra $L^1(G)$. Then $M(L^1(G)) = M(G)$ [Wendel's Theorem]. A bit of measure theory shows that $L^1(G)$ is an ideal in $M(G)$, and so we have an embedding $M(G) \rightarrow M(L^1(G))$. A bounded approximate identity argument gives that this surjects.

Notice that $M(G)$ is always a dual space (and indeed a dual Banach algebra).

Banach algebras; Examples

If A is a Banach algebra, then a little closed graph argument shows that if $(L, R) \in M(A)$, then L and R are bounded. We norm $M(A)$ by regarding it as a subspace of $\mathcal{B}(A) \oplus_{\infty} \mathcal{B}(A)$.

If A is unital, then $A = M(A)$.

If A is a C^* -algebra then so is $M(A)$. For a commutative C^* -algebra $A = C_0(X)$, the multiplier algebra can be identified with $C^b(X)$, which in turn is $C(\beta X)$. So multiplier algebras are Stone-Cech compactifications.

Notice that $M(A)$ is rarely a von Neumann algebra.

Let E be a Banach algebra, and $A = \mathcal{K}(E)$ the compact operators on E . Then $M(A) = \mathcal{B}(E)$.

Notice that $\mathcal{B}(E)$ may or may not be a dual space.

For a locally compact group G , consider the algebra $L^1(G)$. Then $M(L^1(G)) = M(G)$ [Wendel's Theorem]. A bit of measure theory shows that $L^1(G)$ is an ideal in $M(G)$, and so we have an embedding $M(G) \rightarrow M(L^1(G))$. A bounded approximate identity argument gives that this surjects.

Notice that $M(G)$ is always a dual space (and indeed a dual Banach algebra).

Banach algebras; Examples

If A is a Banach algebra, then a little closed graph argument shows that if $(L, R) \in M(A)$, then L and R are bounded. We norm $M(A)$ by regarding it as a subspace of $\mathcal{B}(A) \oplus_{\infty} \mathcal{B}(A)$.

If A is unital, then $A = M(A)$.

If A is a C^* -algebra then so is $M(A)$. For a commutative C^* -algebra $A = C_0(X)$, the multiplier algebra can be identified with $C^b(X)$, which in turn is $C(\beta X)$. So multiplier algebras are Stone-Cech compactifications.

Notice that $M(A)$ is rarely a von Neumann algebra.

Let E be a Banach algebra, and $A = \mathcal{K}(E)$ the compact operators on E . Then $M(A) = \mathcal{B}(E)$.

Notice that $\mathcal{B}(E)$ may or may not be a dual space.

For a locally compact group G , consider the algebra $L^1(G)$. Then $M(L^1(G)) = M(G)$ [Wendel's Theorem]. A bit of measure theory shows that $L^1(G)$ is an ideal in $M(G)$, and so we have an embedding $M(G) \rightarrow M(L^1(G))$. A bounded approximate identity argument gives that this surjects.

Notice that $M(G)$ is always a dual space (and indeed a dual Banach algebra).

Banach algebras; Examples

If A is a Banach algebra, then a little closed graph argument shows that if $(L, R) \in M(A)$, then L and R are bounded. We norm $M(A)$ by regarding it as a subspace of $\mathcal{B}(A) \oplus_{\infty} \mathcal{B}(A)$.

If A is unital, then $A = M(A)$.

If A is a C^* -algebra then so is $M(A)$. For a commutative C^* -algebra $A = C_0(X)$, the multiplier algebra can be identified with $C^b(X)$, which in turn is $C(\beta X)$. So multiplier algebras are Stone-Cech compactifications.

Notice that $M(A)$ is rarely a von Neumann algebra.

Let E be a Banach algebra, and $A = \mathcal{K}(E)$ the compact operators on E . Then $M(A) = \mathcal{B}(E)$.

Notice that $\mathcal{B}(E)$ may or may not be a dual space.

For a locally compact group G , consider the algebra $L^1(G)$. Then $M(L^1(G)) = M(G)$ [Wendel's Theorem]. A bit of measure theory shows that $L^1(G)$ is an ideal in $M(G)$, and so we have an embedding $M(G) \rightarrow M(L^1(G))$. A bounded approximate identity argument gives that this surjects.

Notice that $M(G)$ is always a dual space (and indeed a dual Banach algebra).

Dual Banach algebras

Let A be a Banach algebra which is the dual Banach space of A_* say. We say that A is a dual Banach algebra (for A_*) if the product is separately weak*-continuous.

Let's assemble some ingredients. Let A be a Banach algebra such that $\{ab : a, b \in A\}$ is linearly dense in A . Let (B, B_*) be a dual Banach algebra such that:

- we have an isometric homomorphism $\iota : A \rightarrow B$;
- $\iota(A)$ is an (essential) ideal in B ;
- the resulting map $B \rightarrow M(A)$ injects.

We'll construct a preidual for $M(A)$.

If you are interested in the one-sided case, compare with [Selivanov], Monatsh. Math. (1999).

Dual Banach algebras

Let A be a Banach algebra which is the dual Banach space of A_* say. We say that A is a dual Banach algebra (for A_*) if the product is separately weak*-continuous.

Let's assemble some ingredients. Let A be a Banach algebra such that $\{ab : a, b \in A\}$ is linearly dense in A . Let (B, B_*) be a dual Banach algebra such that:

- we have an isometric homomorphism $\iota : A \rightarrow B$;
- $\iota(A)$ is an (essential) ideal in B ;
- the resulting map $B \rightarrow M(A)$ injects.

We'll construct a predual for $M(A)$.

If you are interested in the one-sided case, compare with [Selivanov], Monatsh. Math. (1999).

Dual Banach algebras

Let A be a Banach algebra which is the dual Banach space of A_* say. We say that A is a dual Banach algebra (for A_*) if the product is separately weak*-continuous.

Let's assemble some ingredients. Let A be a Banach algebra such that $\{ab : a, b \in A\}$ is linearly dense in A . Let (B, B_*) be a dual Banach algebra such that:

- we have an isometric homomorphism $\iota : A \rightarrow B$;
- $\iota(A)$ is an (essential) ideal in B ;
- the resulting map $B \rightarrow M(A)$ injects.

We'll construct a predual for $M(A)$.

If you are interested in the one-sided case, compare with [Selivanov], Monatsh. Math. (1999).

The construction

Consider

$$X = (A \widehat{\otimes} B_*) \oplus_1 (A \widehat{\otimes} B_*) \quad \text{so that} \quad X^* = \mathcal{B}(A, B) \oplus_\infty \mathcal{B}(A, B).$$

Let $Y \subseteq X$ be the linear span of

$$(b \otimes \mu \cdot \iota(a)) \oplus (-a \oplus \iota(b) \cdot \mu) \quad (a, b \in A, \mu \in B_*).$$

Then $Y^\perp \subseteq X^*$ is a weak*-closed subspace with predual X/Y . A calculation shows that

$$(T, S) \in Y^\perp \Leftrightarrow \iota(a)T(b) = S(a)\iota(b) \quad (a, b \in A).$$

Now argue that as products are dense in A , actually $T(A), S(A) \subseteq \iota(A)$, and so we really have maps $L, R : A \rightarrow A$ with $T = \iota L, S = \iota R$. But then $(L, R) \in M(A)$; so we've shown that $M(A) \cong Y^\perp$.

A final, slightly technical, check shows that $M(A)$ is indeed a dual Banach algebra.

The construction

Consider

$$X = (A \widehat{\otimes} B_*) \oplus_1 (A \widehat{\otimes} B_*) \quad \text{so that} \quad X^* = \mathcal{B}(A, B) \oplus_\infty \mathcal{B}(A, B).$$

Let $Y \subseteq X$ be the linear span of

$$(b \otimes \mu \cdot \iota(a)) \oplus (-a \oplus \iota(b) \cdot \mu) \quad (a, b \in A, \mu \in B_*).$$

Then $Y^\perp \subseteq X^*$ is a weak*-closed subspace with predual X/Y . A calculation shows that

$$(T, S) \in Y^\perp \Leftrightarrow \iota(a)T(b) = S(a)\iota(b) \quad (a, b \in A).$$

Now argue that as products are dense in A , actually $T(A), S(A) \subseteq \iota(A)$, and so we really have maps $L, R : A \rightarrow A$ with $T = \iota L, S = \iota R$. But then $(L, R) \in M(A)$; so we've shown that $M(A) \cong Y^\perp$.

A final, slightly technical, check shows that $M(A)$ is indeed a dual Banach algebra.

The construction

Consider

$$X = (A \widehat{\otimes} B_*) \oplus_1 (A \widehat{\otimes} B_*) \quad \text{so that} \quad X^* = \mathcal{B}(A, B) \oplus_\infty \mathcal{B}(A, B).$$

Let $Y \subseteq X$ be the linear span of

$$(b \otimes \mu \cdot \iota(a)) \oplus (-a \oplus \iota(b) \cdot \mu) \quad (a, b \in A, \mu \in B_*).$$

Then $Y^\perp \subseteq X^*$ is a weak*-closed subspace with predual X/Y . A calculation shows that

$$(T, S) \in Y^\perp \Leftrightarrow \iota(a)T(b) = S(a)\iota(b) \quad (a, b \in A).$$

Now argue that as products are dense in A , actually $T(A), S(A) \subseteq \iota(A)$, and so we really have maps $L, R : A \rightarrow A$ with $T = \iota L, S = \iota R$. But then $(L, R) \in M(A)$; so we've shown that $M(A) \cong Y^\perp$.

A final, slightly technical, check shows that $M(A)$ is indeed a dual Banach algebra.

The construction

Consider

$$X = (A \widehat{\otimes} B_*) \oplus_1 (A \widehat{\otimes} B_*) \quad \text{so that} \quad X^* = \mathcal{B}(A, B) \oplus_\infty \mathcal{B}(A, B).$$

Let $Y \subseteq X$ be the linear span of

$$(b \otimes \mu \cdot \iota(a)) \oplus (-a \oplus \iota(b) \cdot \mu) \quad (a, b \in A, \mu \in B_*).$$

Then $Y^\perp \subseteq X^*$ is a weak*-closed subspace with predual X/Y . A calculation shows that

$$(T, S) \in Y^\perp \Leftrightarrow \iota(a)T(b) = S(a)\iota(b) \quad (a, b \in A).$$

Now argue that as products are dense in A , actually $T(A), S(A) \subseteq \iota(A)$, and so we really have maps $L, R : A \rightarrow A$ with $T = \iota L, S = \iota R$. But then $(L, R) \in M(A)$; so we've shown that $M(A) \cong Y^\perp$.

A final, slightly technical, check shows that $M(A)$ is indeed a dual Banach algebra.

Uniqueness

Following the construction through the weak*-topology on $M(A)$ satisfies: a bounded net (L_α, R_α) in $M(A)$ is weak*-null if and only if

$$\lim_{\alpha} \langle \iota L_\alpha(a), \mu \rangle + \langle \iota R_\alpha(b), \lambda \rangle = 0 \quad (a, b \in A, \mu, \lambda \in B_*).$$

Let $\theta : B \rightarrow M(A)$ be the map induced by $\iota : A \rightarrow B$. Then there is one and only one weak*-topology on $M(A)$ such that:

- $M(A)$ is a dual Banach algebra;
- for a bounded net (b_α) in B , we have that (b_α) is weak* null in B if and only if $(\theta(b_\alpha))$ is weak* null in $M(A)$.

So what we've done is taken a dual Banach algebra B which isn't quite large enough to be all of $M(A)$, and boot-strapped the weak*-topology from B to $M(A)$.

Uniqueness

Following the construction through the weak*-topology on $M(A)$ satisfies: a bounded net (L_α, R_α) in $M(A)$ is weak*-null if and only if

$$\lim_{\alpha} \langle \iota L_\alpha(a), \mu \rangle + \langle \iota R_\alpha(b), \lambda \rangle = 0 \quad (a, b \in A, \mu, \lambda \in B_*).$$

Let $\theta : B \rightarrow M(A)$ be the map induced by $\iota : A \rightarrow B$. Then there is one and only one weak*-topology on $M(A)$ such that:

- $M(A)$ is a dual Banach algebra;
- for a bounded net (b_α) in B , we have that (b_α) is weak* null in B if and only if $(\theta(b_\alpha))$ is weak* null in $M(A)$.

So what we've done is taken a dual Banach algebra B which isn't quite large enough to be all of $M(A)$, and boot-strapped the weak*-topology from B to $M(A)$.

Uniqueness

Following the construction through the weak*-topology on $M(A)$ satisfies: a bounded net (L_α, R_α) in $M(A)$ is weak*-null if and only if

$$\lim_{\alpha} \langle \iota L_\alpha(a), \mu \rangle + \langle \iota R_\alpha(b), \lambda \rangle = 0 \quad (a, b \in A, \mu, \lambda \in B_*).$$

Let $\theta : B \rightarrow M(A)$ be the map induced by $\iota : A \rightarrow B$. Then there is one and only one weak*-topology on $M(A)$ such that:

- $M(A)$ is a dual Banach algebra;
- for a bounded net (b_α) in B , we have that (b_α) is weak* null in B if and only if $(\theta(b_\alpha))$ is weak* null in $M(A)$.

So what we've done is taken a dual Banach algebra B which isn't quite large enough to be all of $M(A)$, and boot-strapped the weak*-topology from B to $M(A)$.

The Fourier algebra

Let G be a locally compact group, and let λ be the left-regular representation of G on $L^2(G)$:

$$\lambda(s)\xi : t \mapsto \xi(s^{-1}t) \quad (s, t \in G, \xi \in L^2(G)).$$

Let $VN(G)$ be the group von Neumann algebra, which is generated by $\{\lambda(s) : s \in G\}$.

There exists a normal $*$ -homomorphism $\Delta : VN(G) \rightarrow VN(G \times G)$ which satisfies $\Delta(\lambda(s)) = \lambda(s) \otimes \lambda(s)$. This exists, as we can define a unitary $W \in \mathcal{B}(L^2(G \times G))$ by $W\xi(s, t) = \xi(ts, t)$, and then

$$\Delta(x) = W^*(1 \otimes x)W \quad (x \in VN(G)),$$

does the job.

Let $A(G)$ be the predual of $VN(G)$. As Δ is normal, for $\omega, \sigma \in A(G)$, there exists $\omega\sigma \in A(G)$ such that

$$\langle \Delta(x), \omega \otimes \sigma \rangle = \langle x, \omega\sigma \rangle \quad (x \in VN(G)).$$

Thus we've turned $A(G)$ into a Banach algebra.

The Fourier algebra

Let G be a locally compact group, and let λ be the left-regular representation of G on $L^2(G)$:

$$\lambda(s)\xi : t \mapsto \xi(s^{-1}t) \quad (s, t \in G, \xi \in L^2(G)).$$

Let $VN(G)$ be the group von Neumann algebra, which is generated by $\{\lambda(s) : s \in G\}$.

There exists a normal $*$ -homomorphism $\Delta : VN(G) \rightarrow VN(G \times G)$ which satisfies $\Delta(\lambda(s)) = \lambda(s) \otimes \lambda(s)$. This exists, as we can define a unitary $W \in \mathcal{B}(L^2(G \times G))$ by $W\xi(s, t) = \xi(ts, t)$, and then

$$\Delta(x) = W^*(1 \otimes x)W \quad (x \in VN(G)),$$

does the job.

Let $A(G)$ be the predual of $VN(G)$. As Δ is normal, for $\omega, \sigma \in A(G)$, there exists $\omega\sigma \in A(G)$ such that

$$\langle \Delta(x), \omega \otimes \sigma \rangle = \langle x, \omega\sigma \rangle \quad (x \in VN(G)).$$

Thus we've turned $A(G)$ into a Banach algebra.

The Fourier algebra

Let G be a locally compact group, and let λ be the left-regular representation of G on $L^2(G)$:

$$\lambda(s)\xi : t \mapsto \xi(s^{-1}t) \quad (s, t \in G, \xi \in L^2(G)).$$

Let $VN(G)$ be the group von Neumann algebra, which is generated by $\{\lambda(s) : s \in G\}$.

There exists a normal $*$ -homomorphism $\Delta : VN(G) \rightarrow VN(G \times G)$ which satisfies $\Delta(\lambda(s)) = \lambda(s) \otimes \lambda(s)$. This exists, as we can define a unitary $W \in \mathcal{B}(L^2(G \times G))$ by $W\xi(s, t) = \xi(ts, t)$, and then

$$\Delta(x) = W^*(1 \otimes x)W \quad (x \in VN(G)),$$

does the job.

Let $A(G)$ be the predual of $VN(G)$. As Δ is normal, for $\omega, \sigma \in A(G)$, there exists $\omega\sigma \in A(G)$ such that

$$\langle \Delta(x), \omega \otimes \sigma \rangle = \langle x, \omega\sigma \rangle \quad (x \in VN(G)).$$

Thus we've turned $A(G)$ into a Banach algebra.

As a function algebra

As $\{\lambda(s) : s \in G\}$ generates $VN(G)$, an element $\omega \in A(G)$ is uniquely determined by $\{\langle \lambda(s), \omega \rangle : s \in G\}$ so we can think of ω as a function $G \rightarrow \mathbb{C}; s \mapsto \omega(s) = \langle \lambda(s), \omega \rangle$.

Then the product on $A(G)$ is just the pointwise product, as

$$(\omega\sigma)(s) = \langle \Delta(\lambda(s)), \omega \otimes \sigma \rangle = \langle \lambda(s) \otimes \lambda(s), \omega \otimes \sigma \rangle = \omega(s)\sigma(s).$$

Alternatively, starting with λ , we could integrate this to get a $*$ -homomorphism $\lambda : L^1(G) \rightarrow \mathcal{B}(L^2(G))$. Then the norm closure of the image is $C_r^*(G)$, the (reduced) group C^* -algebra. Then Δ restricts to give a map

$$\Delta : C_r^*(G) \rightarrow M(C_r^*(G \times G)).$$

Using this, we turn $C_r^*(G)^*$ into a commutative, dual Banach algebra, $B_r(G)$ the (reduced) Fourier-Stieltjes algebra.

If G is abelian with dual group \hat{G} , then

$$VN(G) \cong L^\infty(\hat{G}), \quad A(G) \cong L^1(\hat{G}), \quad C_r^*(G) \cong C_0(\hat{G}), \quad B_r(G) \cong M(\hat{G}).$$

As a function algebra

As $\{\lambda(s) : s \in G\}$ generates $VN(G)$, an element $\omega \in A(G)$ is uniquely determined by $\{\langle \lambda(s), \omega \rangle : s \in G\}$ so we can think of ω as a function $G \rightarrow \mathbb{C}; s \mapsto \omega(s) = \langle \lambda(s), \omega \rangle$.

Then the product on $A(G)$ is just the pointwise product, as

$$(\omega\sigma)(s) = \langle \Delta(\lambda(s)), \omega \otimes \sigma \rangle = \langle \lambda(s) \otimes \lambda(s), \omega \otimes \sigma \rangle = \omega(s)\sigma(s).$$

Alternatively, starting with λ , we could integrate this to get a $*$ -homomorphism $\lambda : L^1(G) \rightarrow \mathcal{B}(L^2(G))$. Then the norm closure of the image is $C_r^*(G)$, the (reduced) group C^* -algebra. Then Δ restricts to give a map

$$\Delta : C_r^*(G) \rightarrow M(C_r^*(G \times G)).$$

Using this, we turn $C_r^*(G)^*$ into a commutative, dual Banach algebra, $B_r(G)$ the (reduced) Fourier-Stieltjes algebra.

If G is abelian with dual group \hat{G} , then

$$VN(G) \cong L^\infty(\hat{G}), \quad A(G) \cong L^1(\hat{G}), \quad C_r^*(G) \cong C_0(\hat{G}), \quad B_r(G) \cong M(\hat{G}).$$

As a function algebra

As $\{\lambda(s) : s \in G\}$ generates $VN(G)$, an element $\omega \in A(G)$ is uniquely determined by $\{\langle \lambda(s), \omega \rangle : s \in G\}$ so we can think of ω as a function $G \rightarrow \mathbb{C}; s \mapsto \omega(s) = \langle \lambda(s), \omega \rangle$.

Then the product on $A(G)$ is just the pointwise product, as

$$(\omega\sigma)(s) = \langle \Delta(\lambda(s)), \omega \otimes \sigma \rangle = \langle \lambda(s) \otimes \lambda(s), \omega \otimes \sigma \rangle = \omega(s)\sigma(s).$$

Alternatively, starting with λ , we could integrate this to get a $*$ -homomorphism $\lambda : L^1(G) \rightarrow \mathcal{B}(L^2(G))$. Then the norm closure of the image is $C_r^*(G)$, the (reduced) group C^* -algebra. Then Δ restricts to give a map

$$\Delta : C_r^*(G) \rightarrow M(C_r^*(G \times G)).$$

Using this, we turn $C_r^*(G)^*$ into a commutative, dual Banach algebra, $B_r(G)$ the (reduced) Fourier-Stieltjes algebra.

If G is abelian with dual group \hat{G} , then

$$VN(G) \cong L^\infty(\hat{G}), \quad A(G) \cong L^1(\hat{G}), \quad C_r^*(G) \cong C_0(\hat{G}), \quad B_r(G) \cong M(\hat{G}).$$

As a function algebra

As $\{\lambda(s) : s \in G\}$ generates $VN(G)$, an element $\omega \in A(G)$ is uniquely determined by $\{\langle \lambda(s), \omega \rangle : s \in G\}$ so we can think of ω as a function $G \rightarrow \mathbb{C}; s \mapsto \omega(s) = \langle \lambda(s), \omega \rangle$.

Then the product on $A(G)$ is just the pointwise product, as

$$(\omega\sigma)(s) = \langle \Delta(\lambda(s)), \omega \otimes \sigma \rangle = \langle \lambda(s) \otimes \lambda(s), \omega \otimes \sigma \rangle = \omega(s)\sigma(s).$$

Alternatively, starting with λ , we could integrate this to get a $*$ -homomorphism $\lambda : L^1(G) \rightarrow \mathcal{B}(L^2(G))$. Then the norm closure of the image is $C_r^*(G)$, the (reduced) group C^* -algebra. Then Δ restricts to give a map

$$\Delta : C_r^*(G) \rightarrow M(C_r^*(G \times G)).$$

Using this, we turn $C_r^*(G)^*$ into a commutative, dual Banach algebra, $B_r(G)$ the (reduced) Fourier-Stieltjes algebra.

If G is abelian with dual group \hat{G} , then

$$VN(G) \cong L^\infty(\hat{G}), \quad A(G) \cong L^1(\hat{G}), \quad C_r^*(G) \cong C_0(\hat{G}), \quad B_r(G) \cong M(\hat{G}).$$

Multipliers of $A(G)$

By analogy with the abelian case, we might hope that $M(A(G)) = B_r(G)$ always. However, $B_r(G)$ is unital if and only if G is amenable [Leptin, Cowling?].

As $A(G)$ is commutative, it follows that for $(L, R) \in M(A(G))$ we have $L = R$, and that actually

$$M(A(G)) = \{T : A(G) \rightarrow A(G) : T(\omega\sigma) = T(\omega)\sigma \ (\omega, \sigma \in A(G))\}.$$

Indeed, we can actually identify $M(A(G))$ with a space of functions:

$$M(A(G)) = \{f \in C^b(G) : f\omega \in A(G) \ (\omega \in A(G))\}.$$

We do always have that $A(G) \rightarrow B_r(G) = C_r^*(G)^*$ isometrically (Kaplansky density), and that $A(G)$ is an ideal in $B_r(G)$ (the Fell absorption principle). A little check shows that $A(G)$ is essential in $B_r(G)$. Thus we can run our programme, and $M(A(G))$ is a dual Banach algebra.

This was first shown by [De Canniere, Haagerup]. I think it's nice that we don't really need to know very much about the structure of $A(G)$ at all. Also, this construction happily extends to locally compact quantum groups.

Multipliers of $A(G)$

By analogy with the abelian case, we might hope that $M(A(G)) = B_r(G)$ always. However, $B_r(G)$ is unital if and only if G is amenable [Leptin, Cowling?].

As $A(G)$ is commutative, it follows that for $(L, R) \in M(A(G))$ we have $L = R$, and that actually

$$M(A(G)) = \{T : A(G) \rightarrow A(G) : T(\omega\sigma) = T(\omega)\sigma \ (\omega, \sigma \in A(G))\}.$$

Indeed, we can actually identify $M(A(G))$ with a space of functions:

$$M(A(G)) = \{f \in C^b(G) : f\omega \in A(G) \ (\omega \in A(G))\}.$$

We do always have that $A(G) \rightarrow B_r(G) = C_r^*(G)^*$ isometrically (Kaplansky density), and that $A(G)$ is an ideal in $B_r(G)$ (the Fell absorption principle). A little check shows that $A(G)$ is essential in $B_r(G)$. Thus we can run our programme, and $M(A(G))$ is a dual Banach algebra.

This was first shown by [De Canniere, Haagerup]. I think it's nice that we don't really need to know very much about the structure of $A(G)$ at all. Also, this construction happily extends to locally compact quantum groups.

Multipliers of $A(G)$

By analogy with the abelian case, we might hope that $M(A(G)) = B_r(G)$ always. However, $B_r(G)$ is unital if and only if G is amenable [Leptin, Cowling?].

As $A(G)$ is commutative, it follows that for $(L, R) \in M(A(G))$ we have $L = R$, and that actually

$$M(A(G)) = \{T : A(G) \rightarrow A(G) : T(\omega\sigma) = T(\omega)\sigma \ (\omega, \sigma \in A(G))\}.$$

Indeed, we can actually identify $M(A(G))$ with a space of functions:

$$M(A(G)) = \{f \in C^b(G) : f\omega \in A(G) \ (\omega \in A(G))\}.$$

We do always have that $A(G) \rightarrow B_r(G) = C_r^*(G)^*$ isometrically (Kaplansky density), and that $A(G)$ is an ideal in $B_r(G)$ (the Fell absorption principle). A little check shows that $A(G)$ is essential in $B_r(G)$. Thus we can run our programme, and $M(A(G))$ is a dual Banach algebra.

This was first shown by [De Canniere, Haagerup]. I think it's nice that we don't really need to know very much about the structure of $A(G)$ at all. Also, this construction happily extends to locally compact quantum groups.

Multipliers of $A(G)$

By analogy with the abelian case, we might hope that $M(A(G)) = B_r(G)$ always. However, $B_r(G)$ is unital if and only if G is amenable [Leptin, Cowling?].

As $A(G)$ is commutative, it follows that for $(L, R) \in M(A(G))$ we have $L = R$, and that actually

$$M(A(G)) = \{T : A(G) \rightarrow A(G) : T(\omega\sigma) = T(\omega)\sigma \ (\omega, \sigma \in A(G))\}.$$

Indeed, we can actually identify $M(A(G))$ with a space of functions:

$$M(A(G)) = \{f \in C^b(G) : f\omega \in A(G) \ (\omega \in A(G))\}.$$

We do always have that $A(G) \rightarrow B_r(G) = C_r^*(G)^*$ isometrically (Kaplansky density), and that $A(G)$ is an ideal in $B_r(G)$ (the Fell absorption principle). A little check shows that $A(G)$ is essential in $B_r(G)$. Thus we can run our programme, and $M(A(G))$ is a dual Banach algebra.

This was first shown by [De Canniere, Haagerup]. I think it's nice that we don't really need to know very much about the structure of $A(G)$ at all. Also, this construction happily extends to locally compact quantum groups.

Operator spaces; Completely bounded maps

Given a map $T : A \rightarrow B$ between C^* -algebras, we can dilate T to a map

$$T \otimes \iota_n : A \otimes M_n = M_n(A) \rightarrow B \otimes M_n = M_n(B)$$

between the matrix algebras over A and B . If $\sup_n \|T \otimes \iota_n\| < \infty$, then T is *completely bounded*.

Given a multiplier $T \in M(A(G))$, if $T^* : VN(G) \rightarrow VN(G)$ is completely bounded, then T is completely bounded, $T \in M_{cb}A(G)$.

Of course, the definition of completely bounded makes sense for operators between subspaces of C^* -algebras, and this leads to the notion of an *operator space*. The category of operator spaces and completely bounded maps is nicely behaved, and we can run our construction again, showing that $M_{cb}A(G)$ is a dual Banach algebra.

Operator spaces; Completely bounded maps

Given a map $T : A \rightarrow B$ between C^* -algebras, we can dilate T to a map

$$T \otimes \iota_n : A \otimes M_n = M_n(A) \rightarrow B \otimes M_n = M_n(B)$$

between the matrix algebras over A and B . If $\sup_n \|T \otimes \iota_n\| < \infty$, then T is *completely bounded*.

Given a multiplier $T \in M(A(G))$, if $T^* : VN(G) \rightarrow VN(G)$ is completely bounded, then T is completely bounded, $T \in M_{cb}A(G)$.

Of course, the definition of completely bounded makes sense for operators between subspaces of C^* -algebras, and this leads to the notion of an *operator space*. The category of operator spaces and completely bounded maps is nicely behaved, and we can run our construction again, showing that $M_{cb}A(G)$ is a dual Banach algebra.

Operator spaces; Completely bounded maps

Given a map $T : A \rightarrow B$ between C^* -algebras, we can dilate T to a map

$$T \otimes \iota_n : A \otimes M_n = M_n(A) \rightarrow B \otimes M_n = M_n(B)$$

between the matrix algebras over A and B . If $\sup_n \|T \otimes \iota_n\| < \infty$, then T is *completely bounded*.

Given a multiplier $T \in M(A(G))$, if $T^* : VN(G) \rightarrow VN(G)$ is completely bounded, then T is completely bounded, $T \in M_{cb}A(G)$.

Of course, the definition of completely bounded makes sense for operators between subspaces of C^* -algebras, and this leads to the notion of an *operator space*. The category of operator spaces and completely bounded maps is nicely behaved, and we can run our construction again, showing that $M_{cb}A(G)$ is a dual Banach algebra.

Self-induced algebras and modules

Let $A\widehat{\otimes}_A A$ be the quotient of $A\widehat{\otimes} A$ by the linear closure of

$$\{ab \otimes c - a \otimes bc : a, b, c \in A\}.$$

The product map $\pi : A\widehat{\otimes} A \rightarrow A; a \otimes b \mapsto ab$ respects this quotient, so we get a map $A\widehat{\otimes}_A A \rightarrow A$. If this is an isomorphism, then A is *self-induced*.

Similarly, for a left A -module E , we can form $A\widehat{\otimes}_A E$, and we say that E is *induced* if the product map implements an isomorphism $A\widehat{\otimes}_A E \cong E$.

Let $\theta : A \rightarrow M(B)$ be a homomorphism: we say that this is *non-degenerate* if $\{\theta(a_1)b\theta(a_2) : a_1, a_2 \in A, b \in B\}$ is linearly dense in B . We can turn B into an A -bimodule by setting

$$a \cdot b = \theta(a)b, \quad b \cdot a = b\theta(a) \quad (a \in A, b \in B).$$

Then θ is non-degenerate if B is an essential A -module.

[Johnson] \Rightarrow that if A has a bounded approximate identity, then any non-degenerate homomorphism extends to $\theta : M(A) \rightarrow M(B)$.

Self-induced algebras and modules

Let $A\widehat{\otimes}_A A$ be the quotient of $A\widehat{\otimes} A$ by the linear closure of

$$\{ab \otimes c - a \otimes bc : a, b, c \in A\}.$$

The product map $\pi : A\widehat{\otimes} A \rightarrow A; a \otimes b \mapsto ab$ respects this quotient, so we get a map $A\widehat{\otimes}_A A \rightarrow A$. If this is an isomorphism, then A is *self-induced*.

Similarly, for a left A -module E , we can form $A\widehat{\otimes}_A E$, and we say that E is *induced* if the product map implements an isomorphism $A\widehat{\otimes}_A E \cong E$.

Let $\theta : A \rightarrow M(B)$ be a homomorphism: we say that this is *non-degenerate* if $\{\theta(a_1)b\theta(a_2) : a_1, a_2 \in A, b \in B\}$ is linearly dense in B . We can turn B into an A -bimodule by setting

$$a \cdot b = \theta(a)b, \quad b \cdot a = b\theta(a) \quad (a \in A, b \in B).$$

Then θ is non-degenerate if B is an essential A -module.

[Johnson] \Rightarrow that if A has a bounded approximate identity, then any non-degenerate homomorphism extends to $\theta : M(A) \rightarrow M(B)$.

Self-induced algebras and modules

Let $A\widehat{\otimes}_A A$ be the quotient of $A\widehat{\otimes} A$ by the linear closure of

$$\{ab \otimes c - a \otimes bc : a, b, c \in A\}.$$

The product map $\pi : A\widehat{\otimes} A \rightarrow A; a \otimes b \mapsto ab$ respects this quotient, so we get a map $A\widehat{\otimes}_A A \rightarrow A$. If this is an isomorphism, then A is *self-induced*.

Similarly, for a left A -module E , we can form $A\widehat{\otimes}_A E$, and we say that E is *induced* if the product map implements an isomorphism $A\widehat{\otimes}_A E \cong E$.

Let $\theta : A \rightarrow M(B)$ be a homomorphism: we say that this is *non-degenerate* if $\{\theta(a_1)b\theta(a_2) : a_1, a_2 \in A, b \in B\}$ is linearly dense in B . We can turn B into an A -bimodule by setting

$$a \cdot b = \theta(a)b, \quad b \cdot a = b\theta(a) \quad (a \in A, b \in B).$$

Then θ is non-degenerate if B is an essential A -module.

[Johnson] \Rightarrow that if A has a bounded approximate identity, then any non-degenerate homomorphism extends to $\theta : M(A) \rightarrow M(B)$.

Extending homomorphisms in the induced case

Let $\theta : A \rightarrow M(B)$ be a homomorphism which turns B into an induced A -bimodule. We can extend θ to $M(A)$ as follows. Given $(L, R) \in M(A)$, we define $L' : B \rightarrow B$ by

$$\begin{array}{ccccc}
 B & \xleftarrow{\cong} & A \widehat{\otimes}_A B & \xleftarrow{\quad} & A \widehat{\otimes} B \\
 \downarrow L' & & \downarrow \text{dotted} & & \downarrow L \otimes \iota \\
 B & \xleftarrow{\cong} & A \widehat{\otimes}_A B & \xleftarrow{\quad} & A \widehat{\otimes} B
 \end{array}$$

This commutes, as $L \otimes \iota$ maps $N = \overline{\text{lin}}\{a_1 a_2 \otimes b - a_1 \otimes \theta(a_2) b : a_1, a_2 \in A, b \in B\}$ into itself. This follows, as

$$L(a_1 a_2) \otimes b - L(a_1) \otimes \theta(a_2) b = L(a_1) a_2 \otimes b - L(a_1) \otimes \theta(a_2) b \in N.$$

Similarly we form $R' : B \rightarrow B$, check that $(L', R') \in M(A)$, and that $(L, R) \mapsto (L', R')$ is homomorphism.

Example of an induced algebra

[Rieffel; Forrest, Lee, Samei] \Rightarrow If A has a bounded approximate identity, then A is self-induced and any essential module is induced.

We now work in the category of Operator Spaces. We'll show that $A(G)$ is always self-induced: we want that $A(G) \widehat{\otimes} A(G)/N = A(G \times G)/N$ is isomorphic to $A(G)$ under the product map. Dualising, we want that

$$VN(G) \xrightarrow{\Delta} N^\perp \subseteq VN(G) \overline{\otimes} VN(G) = VN(G \times G)$$

is an isomorphism. All we need to prove is that it's onto. However,

$$\begin{aligned} N^\perp &= \{x \in VN(G) \overline{\otimes} VN(G) : \langle x, ab \otimes c - a \otimes bc \rangle = 0\} \\ &= \{x \in VN(G) \overline{\otimes} VN(G) : \langle (\Delta \otimes \iota)x - (\iota \otimes \Delta)x, a \otimes b \otimes c \rangle = 0\} \\ &= \{x \in VN(G) \overline{\otimes} VN(G) : (\Delta \otimes \iota)x = (\iota \otimes \Delta)x\}. \end{aligned}$$

We can then check that the *support* of $x \in N^\perp$ must be contained in the diagonal $\{(s, s) : s \in G\} \subseteq G \times G$. As this is a set of synthesis, by [Herz; Takesaki, Tatsumma] we have that $x \in \Delta(VN(G))$, so we're done.

Example of an induced algebra

[Rieffel; Forrest, Lee, Samei] \Rightarrow If A has a bounded approximate identity, then A is self-induced and any essential module is induced.

We now work in the category of Operator Spaces. We'll show that $A(G)$ is always self-induced: we want that $A(G) \widehat{\otimes} A(G)/N = A(G \times G)/N$ is isomorphic to $A(G)$ under the product map. Dualising, we want that

$$VN(G) \xrightarrow{\Delta} N^\perp \subseteq VN(G) \overline{\otimes} VN(G) = VN(G \times G)$$

is an isomorphism. All we need to prove is that it's onto. However,

$$\begin{aligned} N^\perp &= \{x \in VN(G) \overline{\otimes} VN(G) : \langle x, ab \otimes c - a \otimes bc \rangle = 0\} \\ &= \{x \in VN(G) \overline{\otimes} VN(G) : \langle (\Delta \otimes \iota)x - (\iota \otimes \Delta)x, a \otimes b \otimes c \rangle = 0\} \\ &= \{x \in VN(G) \overline{\otimes} VN(G) : (\Delta \otimes \iota)x = (\iota \otimes \Delta)x\}. \end{aligned}$$

We can then check that the *support* of $x \in N^\perp$ must be contained in the diagonal $\{(s, s) : s \in G\} \subseteq G \times G$. As this is a set of synthesis, by [Herz; Takesaki, Tatsumma] we have that $x \in \Delta(VN(G))$, so we're done.

Example of an induced algebra

[Rieffel; Forrest, Lee, Samei] \Rightarrow If A has a bounded approximate identity, then A is self-induced and any essential module is induced.

We now work in the category of Operator Spaces. We'll show that $A(G)$ is always self-induced: we want that $A(G) \widehat{\otimes} A(G)/N = A(G \times G)/N$ is isomorphic to $A(G)$ under the product map. Dualising, we want that

$$VN(G) \xrightarrow{\Delta} N^\perp \subseteq VN(G) \overline{\otimes} VN(G) = VN(G \times G)$$

is an isomorphism. All we need to prove is that it's onto. However,

$$\begin{aligned} N^\perp &= \{x \in VN(G) \overline{\otimes} VN(G) : \langle x, ab \otimes c - a \otimes bc \rangle = 0\} \\ &= \{x \in VN(G) \overline{\otimes} VN(G) : \langle (\Delta \otimes \iota)x - (\iota \otimes \Delta)x, a \otimes b \otimes c \rangle = 0\} \\ &= \{x \in VN(G) \overline{\otimes} VN(G) : (\Delta \otimes \iota)x = (\iota \otimes \Delta)x\}. \end{aligned}$$

We can then check that the *support* of $x \in N^\perp$ must be contained in the diagonal $\{(s, s) : s \in G\} \subseteq G \times G$. As this is a set of synthesis, by [Herz; Takesaki, Tatsumma] we have that $x \in \Delta(VN(G))$, so we're done.

Other extensions

For a homomorphism $\theta : A \rightarrow M(B)$, we can extend to $M(A)$ when:

- A has a bounded approximate identity, and θ is non-degenerate;
- B becomes an induced A -bimodule;
- [Ilie, Stokke] A has a bounded approximate identity, and $M(B)$ is a dual Banach algebra.

For more on induced algebras, see work of [Gronbaek].

Haagerup tensor products

For a C^* -algebra A , we define a norm $\|\cdot\|_h$ on $A \otimes A$ by

$$\|\tau\|_h = \inf \left\{ \left\| \sum_k a_k a_k^* \right\| \left\| \sum_k b_k^* b_k \right\| : \tau = \sum_k a_k \otimes b_k \right\}.$$

Write $A \otimes^h A$ for the completed tensor product.

If $A \subseteq \mathcal{B}(H)$ then the (maybe infinite) column vector $b = (b_1, b_2, \dots)^T$ can be regarded as a map $H \rightarrow H^{(\infty)}$, and then $b^* b = \sum_k b_k^* b_k$. Similarly the row vector $a = (a_1, a_2, \dots)$ is such that $aa^* = \sum_k a_k a_k^*$. Then notice that

$$ab = \sum_k a_k b_k, \quad \|ab\| \leq \|a\| \|b\| = \|aa^*\|^{1/2} \|b^* b\|^{1/2} = \left\| \sum_k a_k a_k^* \right\| \left\| \sum_k b_k^* b_k \right\|.$$

So multiplication $A \otimes^h A \rightarrow A$; $a \otimes b \mapsto ab$ is (completely) contractive. (Multiplication from $A \otimes_{\min} A$ is rarely even bounded).

Haagerup tensor products

For a C^* -algebra A , we define a norm $\|\cdot\|_h$ on $A \otimes A$ by

$$\|\tau\|_h = \inf \left\{ \left\| \sum_k a_k a_k^* \right\| \left\| \sum_k b_k^* b_k \right\| : \tau = \sum_k a_k \otimes b_k \right\}.$$

Write $A \otimes^h A$ for the completed tensor product.

If $A \subseteq \mathcal{B}(H)$ then the (maybe infinite) column vector $b = (b_1, b_2, \dots)^T$ can be regarded as a map $H \rightarrow H^{(\infty)}$, and then $b^* b = \sum_k b_k^* b_k$. Similarly the row vector $a = (a_1, a_2, \dots)$ is such that $aa^* = \sum_k a_k a_k^*$. Then notice that

$$ab = \sum_k a_k b_k, \quad \|ab\| \leq \|a\| \|b\| = \|aa^*\|^{1/2} \|b^* b\|^{1/2} = \left\| \sum_k a_k a_k^* \right\| \left\| \sum_k b_k^* b_k \right\|.$$

So multiplication $A \otimes^h A \rightarrow A$; $a \otimes b \mapsto ab$ is (completely) contractive. (Multiplication from $A \otimes_{\min} A$ is rarely even bounded).

Haagerup tensor products (cont.)

The same definition works to form $A \otimes^h B$. The Haagerup tensor product is injective (and projective). So it makes perfect sense on Operator Spaces as well. The Haagerup tensor product is then *self-dual* in the sense that

$$E^* \otimes^h F^* \subseteq (E \otimes^h F)^* \text{ (completely) isometrically,}$$

for Operator Spaces E and F .

For a von Neumann algebra M , we define the *extended Haagerup tensor product* by

$$M \otimes^{eh} M = (M_* \otimes^h M_*)^*.$$

(This can also be defined as before, but with the sums $\sum_k a_k a_k^*$, and so forth, being interpreted in the σ -weak topology, not the norm topology).

For an operator space E , we define

$$E \otimes^{eh} E = (E^* \otimes^h E^*)_{\sigma}^*,$$

the separately-weak*-continuous functionals $E^* \otimes^h E^* \rightarrow \mathbb{C}$.

Hopf convolution algebras

I introduced the Fourier algebra by:

- defining a von Neumann algebra $VN(G)$;
- defining a “coproduct”, a normal $*$ -homomorphism $\Delta : VN(G) \rightarrow VN(G) \overline{\otimes} VN(G)$ with $(\Delta \otimes \iota)\Delta = (\iota \otimes \Delta)\Delta$.
- then Δ induces the algebra structure on $A(G) = VN(G)_*$.

Can we do everything at the level of $A(G)$? We'd need a map

$$m : A(G) \rightarrow A(G) \otimes A(G) \quad \text{for some tensor product}$$

whose adjoint $m : VN(G) \otimes VN(G) \rightarrow VN(G)$ was the product on $VN(G)$.

[Quigg] tried to do with with the projective tensor product, but that only works (in any sense) if G is abelian by finite.

We could try the Haagerup tensor product $A(G) \otimes^h A(G)$, as then the adjoint is the product map $VN(G) \otimes^{eh} VN(G) \rightarrow VN(G)$, which is (completely) contractive.

Hopf convolution algebras

I introduced the Fourier algebra by:

- defining a von Neumann algebra $VN(G)$;
- defining a “coproduct”, a normal $*$ -homomorphism $\Delta : VN(G) \rightarrow VN(G) \overline{\otimes} VN(G)$ with $(\Delta \otimes \iota)\Delta = (\iota \otimes \Delta)\Delta$.
- then Δ induces the algebra structure on $A(G) = VN(G)_*$.

Can we do everything at the level of $A(G)$? We'd need a map

$$m : A(G) \rightarrow A(G) \otimes A(G) \quad \text{for some tensor product}$$

whose adjoint $m : VN(G) \otimes VN(G) \rightarrow VN(G)$ was the product on $VN(G)$.

[Quigg] tried to do with with the projective tensor product, but that only works (in any sense) if G is abelian by finite.

We could try the Haagerup tensor product $A(G) \otimes^h A(G)$, as then the adjoint is the product map $VN(G) \otimes^{eh} VN(G) \rightarrow VN(G)$, which is (completely) contractive.

The problem

However, this is too small: even if G is abelian but non-compact, then $A(G) \otimes^h A(G)$ won't work.

[Effros+Ruan] solve this by working with $A(G) \otimes^{eh} A(G)$. So we get a map $m : A(G) \rightarrow A(G) \otimes^{eh} A(G)$. For a good analogy, this should be a homomorphism.

To turn $A(G) \otimes^{eh} A(G)$ into an algebra, we use the *shuffle map*

$$S : (A(G) \otimes^{eh} A(G)) \widehat{\otimes} (A(G) \otimes^{eh} A(G)) \rightarrow (A(G) \widehat{\otimes} A(G)) \otimes^{eh} (A(G) \widehat{\otimes} A(G))$$

which sends $(a \otimes b) \otimes (c \otimes d)$ to $(a \otimes c) \otimes (b \otimes d)$. Then the product on $A(G) \otimes^{eh} A(G)$ is given by:

$$\begin{array}{ccc} (A(G) \otimes^{eh} A(G)) \widehat{\otimes} (A(G) \otimes^{eh} A(G)) & \xrightarrow{S} & (A(G) \widehat{\otimes} A(G)) \otimes^{eh} (A(G) \widehat{\otimes} A(G)) \\ & & \downarrow \Delta_* \otimes \Delta_* \\ & & A(G) \otimes^{eh} A(G) \end{array}$$

The problem

However, this is too small: even if G is abelian but non-compact, then $A(G) \otimes^h A(G)$ won't work.

[Effros+Ruan] solve this by working with $A(G) \otimes^{eh} A(G)$. So we get a map $m : A(G) \rightarrow A(G) \otimes^{eh} A(G)$. For a good analogy, this should be a homomorphism.

To turn $A(G) \otimes^{eh} A(G)$ into an algebra, we use the *shuffle map*

$$S : (A(G) \otimes^{eh} A(G)) \widehat{\otimes} (A(G) \otimes^{eh} A(G)) \rightarrow (A(G) \widehat{\otimes} A(G)) \otimes^{eh} (A(G) \widehat{\otimes} A(G))$$

which sends $(a \otimes b) \otimes (c \otimes d)$ to $(a \otimes c) \otimes (b \otimes d)$. Then the product on $A(G) \otimes^{eh} A(G)$ is given by:

$$\begin{array}{ccc} (A(G) \otimes^{eh} A(G)) \widehat{\otimes} (A(G) \otimes^{eh} A(G)) & \xrightarrow{S} & (A(G) \widehat{\otimes} A(G)) \otimes^{eh} (A(G) \widehat{\otimes} A(G)) \\ & & \downarrow \Delta_* \otimes \Delta_* \\ & & A(G) \otimes^{eh} A(G) \end{array}$$

Using multipliers

Remember that, if I don't like von Neumann algebras, I can work with $C_r^*(G)$. However, here Δ now maps from $C_r^*(G)$ to the multiplier algebra $M(C_r^*(G) \otimes_{\min} C_r^*(G))$.

So can we analogously replace $A(G) \otimes^{eh} A(G)$ by, for example, $M_{cb}(A(G) \otimes^h A(G))$? (So, if G is compact, then actually we should be able to work with $A(G) \otimes^h A(G)$).

We can indeed do so. Firstly note that $A(G) \otimes^h A(G) \subseteq A(G) \otimes^{eh} A(G)$ (completely) isometrically, and so it follows that $A(G) \otimes^h A(G)$ is a subalgebra of $A(G) \otimes^{eh} A(G)$. The strategy will be to show that $m : A(G) \rightarrow A(G) \otimes^{eh} A(G)$ maps into the *idealiser*

$$\{\tau \in A(G) \otimes^{eh} A(G) : \tau\sigma \in A(G) \otimes^h A(G) \ (\sigma \in A(G) \otimes^h A(G))\}.$$

If we can do this, then we obviously have a map $A(G) \rightarrow M_{cb}(A(G) \otimes^h A(G))$.

Using multipliers

Remember that, if I don't like von Neumann algebras, I can work with $C_r^*(G)$. However, here Δ now maps from $C_r^*(G)$ to the multiplier algebra $M(C_r^*(G) \otimes_{\min} C_r^*(G))$.

So can we analogously replace $A(G) \otimes^{eh} A(G)$ by, for example, $M_{cb}(A(G) \otimes^h A(G))$? (So, if G is compact, then actually we should be able to work with $A(G) \otimes^h A(G)$).

We can indeed do so. Firstly note that $A(G) \otimes^h A(G) \subseteq A(G) \otimes^{eh} A(G)$ (completely) isometrically, and so it follows that $A(G) \otimes^h A(G)$ is a subalgebra of $A(G) \otimes^{eh} A(G)$. The strategy will be to show that $m : A(G) \rightarrow A(G) \otimes^{eh} A(G)$ maps into the *idealiser*

$$\{\tau \in A(G) \otimes^{eh} A(G) : \tau\sigma \in A(G) \otimes^h A(G) \ (\sigma \in A(G) \otimes^h A(G))\}.$$

If we can do this, then we obviously have a map $A(G) \rightarrow M_{cb}(A(G) \otimes^h A(G))$.

The proof

An alternative description of $(E \otimes^h F)^*$ is those maps $F \rightarrow E^*$ which cb-factor through a *column* Hilbert space. For $a, b, c \in A(G)$, consider $m(a)(b \otimes c) \in A(G) \otimes^{eh} A(G) = (VN(G) \otimes^h VN(G))_{\sigma}^*$; let's see how to view this as a map which factors through the column $L^2(G) = \mathcal{B}(\mathbb{C}, L^2(G))$:

$$\begin{array}{ccccc} VN(G) & \xrightarrow{\hspace{10em}} & A(G) & \hookrightarrow & VN(G)^* \\ & \searrow \alpha & & \nearrow \beta & \\ & & \mathcal{B}(\mathbb{C}, L^2(G)) & & \end{array}$$

Let $a \in A(G)$ be the normal functional $\langle x, a \rangle = (x\xi_0 | \eta_0)$ for $x \in VN(G)$. Then we have

$$\begin{aligned} \alpha(x) &= (c \cdot x)(\xi_0) & (x \in VN(G)), \\ \beta(\xi) &= \omega_{\xi, \eta_0} b & (\xi \in L^2(G)), \end{aligned}$$

where $\omega_{\xi, \eta_0} : VN(G) \rightarrow \mathbb{C}; x \mapsto (x\xi | \eta_0)$.

The proof continued

Let's think more about $\alpha : VN(G) \rightarrow L^2(G); x \mapsto (c \cdot x)(\xi_0)$. Let $c = \omega_{\xi_1, \eta_1}$, so that

$$(\alpha(x)|\eta) = \langle c \cdot x, \omega_{\xi_0, \eta} \rangle = (\Delta(x)\xi_0 \otimes \xi_2 | \eta \otimes \eta_2) = ((1 \otimes x)W(\xi_0 \otimes \xi_2) | W(\eta \otimes \eta_2)).$$

Let (e_i) be an orthonormal basis for $L^2(G)$, so we can write

$$W(\xi_0 \otimes \xi_2) = \sum_i e_i \otimes \phi_i,$$

for some (ϕ_i) . Then, letting $\sigma : L^2(G) \otimes L^2(G) \rightarrow L^2(G) \otimes L^2(G)$ be the swap map,

$$(\alpha(x)|\eta) = \sum_i (\sigma W^*(e_i \otimes x(\phi_i)) | \eta_2 \otimes \eta) = \sum_i ((\omega_{e_i, \eta_2} \otimes \iota)(\sigma W^*)x\phi_i | \eta).$$

Now, the key idea is that $(\omega_{e_i, \eta_2} \otimes \iota)(\sigma W^*)$ is a compact operator, which we can approximate by finite-ranks. So, being careful, we can approximate α , in the cb-norm, by a finite-rank map.

Finishing up

Recall we had $a, b, c \in A(G)$, and we viewed $m(a)(b \otimes c)$ as:

$$\begin{array}{ccccc} VN(G) & \xrightarrow{\quad} & A(G) & \hookrightarrow & VN(G)^* \\ & \searrow \alpha & & \nearrow \beta & \\ & & B(\mathbb{C}, L^2(G)) & & \end{array}$$

We can cb-norm approximate α by a finite-rank map, so we can approximate $m(a)(b \otimes c)$, in the extended Haagerup norm, by a finite-rank tensor in $A(G) \otimes A(G)$. As $A(G) \otimes^h A(G)$ is closed in $A(G) \otimes^{eh} A(G)$, it follows that $m(a)(b \otimes c) \in A(G) \otimes^h A(G)$, as required.

Various open problems

We have a complete contraction $A(G) \rightarrow C_0(G)$ and so also complete contractions

$$\begin{aligned} A(G \times G) &= A(G) \widehat{\otimes} A(G) \rightarrow A(G) \otimes^h A(G) \rightarrow C_0(G) \otimes^h C_0(G) \\ &\rightarrow C_0(G) \otimes_{\min} C_0(G) = C_0(G \times G). \end{aligned}$$

So we can view $\mathfrak{A} = A(G) \otimes^h A(G)$ as an algebra of functions on $G \times G$. A result of [Gelbaum, Robbins] easily generalises to show that the spectrum of \mathfrak{A} is $G \times G$.

- What sort of spectral synthesis properties does \mathfrak{A} have?
- When is $m : A(G) \rightarrow M_{cb}(A(G) \otimes^h A(G))$ induced?
- Can we otherwise form an extension $M_{cb}A(G) \rightarrow M_{cb}(A(G) \otimes^h A(G))$ when G is non-amenable?

Is there any use of the fact that $A(G)$ is a (completely contractive) self-induced algebra? For a general locally compact quantum group, do we have $\{x \in L^\infty(\mathbb{G}) : (\Delta \otimes \iota)x = (\iota \otimes \Delta)x\} = \Delta(L^\infty(\mathbb{G}))$?