

Perspectives on Noncommutative Graphs


Matthew Daws

UCLan

Seoul National University, 19th April 2023

Graphs

A graph consists of a (finite) set of *vertices* V and a collection of *edges* $E \subseteq V \times V$.



$$V = \{A, B, C\} \text{ say, and } E = \{(A, B), (B, C), (C, B), (C, A)\}.$$

A graph is *undirected* if $(x, y) \in E \Leftrightarrow (y, x) \in E$. We allow *self-loops*, so $(x, x) \in E$.

Notice that a graph $G = (V, E)$ is exactly a *relation* on the set V . An undirected graph gives a symmetric relation; having a loop on each vertex gives a reflexive relation.

Channels

A channel sends an input message (element of a finite set A) to an output message (element of a finite set B) perhaps with *noise* so that there is a probability that $a \in A$ is mapped to different $b \in B$.

- Input “o” might be sent to “o” or “0” or “a”.

$p(b|a)$ = probability that b is received given that a was sent

Define a (simple, undirected) graph structure on A by

(a_1, a_2) an edge when $p(b|a_1)p(b|a_2) > 0$ for some b .

This is the *confusability graph* of the channel.

If we want to communicate with *zero error* then we seek a maximal *independent set* in A .

Quantum Mechanics

- A *state* is a unit vector $|\psi\rangle$ in a (finite dim) Hilbert space H .
- More generally, a *density* is a positive, trace-one operator $\rho \in \mathcal{B}(H)$.
- A rank-one density is always of the form $|\psi\rangle\langle\psi|$ for some state ψ .
- (Use Trace duality, so $\omega \in \mathcal{B}(H)^*$ is associated uniquely to $A \in \mathcal{B}(H)$ with $\omega(T) = \text{tr}(AT)$. Then densities are exactly the *states* on $\mathcal{B}(H)$. Here we “overload” the term “state”!)

A (*quantum*) *channel* is a trace-preserving, completely positive (CPTP) map $\mathcal{B}(H_A) \rightarrow \mathcal{B}(H_B)$:

- positive and trace-preserving so it maps densities to densities;
- completely positive so you can tensor with another system and still have positivity.

Stinespring and Kraus

The Stinespring Representation Theorem tells us that any CP map $\mathcal{E} : \mathcal{B}(H_A) \rightarrow \mathcal{B}(H_B)$ has the form

$$\mathcal{E}(x) = V^* \pi(x) V \quad (x \in \mathcal{B}(H_A)),$$

where $V : H_B \rightarrow K$, and $\pi : \mathcal{B}(H_A) \rightarrow \mathcal{B}(K)$ is a $*$ -representation.

- Any such π is of the form $\pi(x) = x \otimes 1$ where $K \cong H_A \otimes K'$.
- Take an o.n. basis (e_i) for K' , so $V(\xi) = \sum_i K_i^*(\xi) \otimes e_i$ for some operators $K_i : H_A \rightarrow H_B$.

We arrive at the *Kraus form*:

$$\mathcal{E}(x) = \sum_i K_i x K_i^* \quad (x \in \mathcal{B}(H_A)).$$

Trace-preserving when $\sum_i K_i^* K_i = 1$.

Quantum zero-error

We turn $\mathcal{B}(H)$ into a Hilbert space using the trace: $(T|S) = \text{tr}(T^*S)$. We say that we can *distinguish* densities when they are orthogonal. Let $\mathcal{E}(x) = \sum_i K_i x K_i^*$ be a quantum channel. We ask when we can distinguish $\mathcal{E}(\rho)$ and $\mathcal{E}(\sigma)$, i.e. when $\mathcal{E}(\rho) \perp \mathcal{E}(\sigma)$. As \mathcal{E} is positive, this is equivalent to

$$\mathcal{E}(|\psi\rangle\langle\psi|) \perp \mathcal{E}(|\phi\rangle\langle\phi|) \quad (\psi \in \text{Im } \rho, \phi \in \text{Im } \sigma).$$

Thus

$$\begin{aligned} 0 &= \text{tr}(\mathcal{E}(|\psi\rangle\langle\psi|)\mathcal{E}(|\phi\rangle\langle\phi|)) = \sum_{i,j} \text{tr}(K_i|\psi\rangle\langle\psi|K_i^* K_j|\phi\rangle\langle\phi|K_j^*) \\ &= \sum_{i,j} |\langle\psi|K_i^* K_j|\phi\rangle|^2 \end{aligned}$$

which is equivalent to $\langle\psi|K_i^* K_j|\phi\rangle = 0$ for each i, j .

To operator systems

So ψ, ϕ are distinguishable after applying \mathcal{E} when

$$\langle \psi | T | \phi \rangle = 0 \quad \text{for each } T \in \text{lin}\{K_i^* K_j\}.$$

Set $\mathcal{S} = \text{lin}\{K_i^* K_j\}$ which has properties:

- \mathcal{S} is a linear subspace;
- $T \in \mathcal{S}$ if and only if $T^* \in \mathcal{S}$;
- $1 \in \mathcal{S}$ (as $\sum_i K_i^* K_i = 1$ as \mathcal{E} is CPTP).

That is, \mathcal{S} is an *operator system*, which depends only on \mathcal{E} and not the choice of (K_i) .

Theorem (Duan)

For any operator system $\mathcal{S} \subseteq \mathcal{B}(H_A)$ there is some quantum channel $\mathcal{E} : \mathcal{B}(H_A) \rightarrow \mathcal{B}(H_B)$ giving rise to \mathcal{S} .

In the classical case

Given a classical channel from A to B with probabilities $p(b|a)$, define Kraus operators

$$K_{ab} = p(b|a)^{1/2} |b\rangle\langle a| : H_A \rightarrow H_B.$$

Here $(|a\rangle)$ is the canonical basis of $H_A = \ell^2(A) \cong \mathbb{C}^{|A|}$.

$$\sum_{ab} K_{ab} |c\rangle\langle c| K_{ab}^* = \sum_{ab} p(b|a) |b\rangle\langle a|c\rangle\langle c|a\rangle\langle b| = \sum_b p(b|c) |b\rangle\langle b|.$$

So the pure state $|c\rangle\langle c|$ is mapped to the combination of pure states which can be received, given that message c is sent.

$$\begin{aligned} \mathcal{S} &= \text{lin}\{K_{ab}^* K_{cd}\} = \text{lin}\{p(b|a)^{1/2} p(d|c)^{1/2} |a\rangle\langle b|d\rangle\langle c|\} \\ &= \text{lin}\{|a\rangle\langle c| : a \sim c\} \end{aligned}$$

Thus \mathcal{S} is directly linked to the confusability graph of the channel.

Quantum relations

Simultaneously, and motivated more by “noncommutative geometry”, Weaver studied:

Definition

Let $M \subseteq \mathcal{B}(H)$ be a von Neumann algebra. A *quantum relation* on M is a weak*-closed subspace $S \subseteq \mathcal{B}(H)$ with $M'SM' \subseteq S$. We define the relation to be:

- 1 *reflexive* when $M' \subseteq S$;
- 2 *symmetric* when $S^* = S$ where $S^* = \{x^* : x \in S\}$;
- 3 *transitive* when $S^2 \subseteq S$, where $S^2 = \overline{\text{lin}}^{w^*} \{xy : x, y \in S\}$.

When $M = \ell^\infty(X) \subseteq \mathcal{B}(\ell^2(X))$ there is a bijection between the usual meaning of “relation” on X and quantum relations on M , given by

$$S = \overline{\text{lin}}^{w^*} \{e_{x,y} : x \sim y\}.$$

Quantum graphs

As a graph on a (finite) vertex set V is simply a relation, and

- undirected graph corresponds to a symmetric relation;
- a reflexive relation corresponds to having a “loop” at every vertex.

Definition (Weaver)

A *quantum graph* on a von Neumann algebra $M \subseteq \mathcal{B}(H)$ is a reflexive, symmetric quantum relation. That is, a unital, self-adjoint, weak*-closed subspace $S \subseteq \mathcal{B}(H)$, which is an M' -bimodule ($M'SM' \subseteq S$).

If $M = \mathcal{B}(H)$ with H finite-dimensional, then as $M' = \mathbb{C}$, a quantum graph is just an operator system: that is, exactly what we had before!
[Duan, Severini, Winter; Stahlke]

Adjacency matrices

Given a graph $G = (V, E)$ consider the $\{0, 1\}$ -valued matrix A with

$$A_{i,j} = \begin{cases} 1 & : (i, j) \in E, \\ 0 & : \text{otherwise,} \end{cases}$$

the *adjacency matrix* of G .

- A is idempotent for the *Schur product*;
- G is undirected if and only if A is self-adjoint;
- A has 1s down the diagonal when G has a loop at every vertex.

We can think of A as an operator on $\ell^2(V)$. This is the GNS space for the C^* -algebra $\ell^\infty(V)$ for the state induced by the uniform measure.

General C^* -algebras

Let B be a finite-dimensional C^* -algebra, and let φ be a faithful state on B , with GNS space $L^2(B)$. Thus B bijects with $L^2(B)$ as a vector space, and so we get:

- The multiplication on B induces a map $m : L^2(B) \otimes L^2(B) \rightarrow L^2(B)$;
- The unit in B induces a map $\eta : \mathbb{C} \rightarrow L^2(B)$.

We get an analogue of the Schur product:

$$x \bullet y = m(x \otimes y)m^* \quad (x, y \in \mathcal{B}(L^2(B))).$$

Quantum adjacency matrix

Definition (Many authors)

A *quantum adjacency matrix* is a self-adjoint $A \in \mathcal{B}(L^2(B))$ with:

- $m(A \otimes A)m^* = A$ (so Schur product idempotent);
- $(1 \otimes \eta^* m)(1 \otimes A \otimes 1)(m^* \eta \otimes 1) = A$;
- $m(A \otimes 1)m^* = \text{id}$ (a “loop at every vertex”);

The middle axiom is a little mysterious: it roughly corresponds to “undirected”.

I want to sketch why this definition is equivalent to the previous notion of a “quantum graph”.

Subspaces to projections

Fix a finite-dimensional C^* -algebra (von Neumann algebra) M . A “quantum graph” is either:

- A subspace of $\mathcal{B}(H)$ (where $M \subseteq \mathcal{B}(H)$) with some properties; or
- An operator on $L^2(M)$ with some properties.

How do we move between these?

$S \subseteq \mathcal{B}(H)$ is a bimodule over M' . As H is finite-dimensional, $\mathcal{B}(H)$ is a Hilbert space for

$$(x|y) = \text{tr}(x^*y).$$

Then $M \otimes M^{\text{op}}$ is represented on $\mathcal{B}(H)$ via



$$\pi : M \otimes M^{\text{op}} \rightarrow \mathcal{B}(\mathcal{B}(H)); \quad \pi(x \otimes y) : T \mapsto xTy.$$

- The commutant of $\pi(M \otimes M^{\text{op}})$ is naturally $M' \otimes (M')^{\text{op}}$.
- So an M' -bimodule of $\mathcal{B}(H)$ corresponds to an $M' \otimes (M')^{\text{op}}$ -invariant subspace of the Hilbert space $\mathcal{B}(H)$;
- Which corresponds to a *projection* in $M \otimes M^{\text{op}}$.

Operators to algebras

So how can we relate:

- Operators $A \in \mathcal{B}(L^2(M))$ with
- Projections in $M \otimes M^{\text{op}}$?



[Musto, Reutter, Verdon]

Operators to algebras 2

Recall the GNS construction for a *tracial* state ψ on M :

$$\Lambda : M \rightarrow L^2(M); \quad (\Lambda(x)|\Lambda(y)) = \psi(x^*y).$$

As $L^2(M)$ is finite-dimensional, every operator on $L^2(M)$ is a linear combination of rank-one operators, and each rank-one operator is of the form

$$\theta_{\Lambda(a),\Lambda(b)} : \xi \mapsto (\Lambda(a)|\xi)\Lambda(b) \quad (\xi \in L^2(M)).$$

Define a bijection

$$\Psi : \mathcal{B}(L^2(M)) \rightarrow M \otimes M^{\text{op}}; \quad \theta_{\Lambda(a),\Lambda(b)} = b \otimes a^*,$$

and extend by linearity!

Operators to algebras 3

$$\Psi : \mathcal{B}(L^2(M)) \rightarrow M \otimes M^{\text{op}}; \quad \theta_{\wedge(a), \wedge(b)} = b \otimes a^*,$$

- Ψ is a homomorphism for the “Schur product”
 $A_1 \bullet A_2 = m(A_1 \otimes A_2)m^*$;
- $A \mapsto (1 \otimes \eta^* m)(1 \otimes A \otimes 1)(m^* \eta \otimes 1)$ corresponds to the anti-homomorphism $\sigma : a \otimes b \mapsto b \otimes a$;
- $A \mapsto A^*$ corresponds to $e \mapsto \sigma(e)^*$.

Conclude: A quantum adjacency matrix corresponds to a projection e with $\sigma(e) = e$. BUT: There is no clean one-to-one correspondence between the axioms.

Non-tracial case

If the functional ψ on M is not tracial, then this correspondence fails.
However:

Theorem (D.)

There is a bijection between:

- “Schur idempotent”, self-adjoint operators A on $L^2(M)$;
- $e \in M \otimes M^{\text{op}}$ with $e^2 = e$ and $e = \sigma(e)^*$;
- self-adjoint M' -bimodules $S \subseteq \mathcal{B}(H)$ such that there is another self-adjoint M' -bimodule S_0 with $S \oplus S_0 = \mathcal{B}(H)$

KMS States

Any faithful state ψ is KMS: there is an automorphism σ' of M with

$$\psi(ab) = \psi(b\sigma'(a)) \quad (a, b \in M).$$

Indeed, there is $Q \in M$ positive and invertible with

$$\psi(a) = \text{tr}(Qa) \quad \text{and then} \quad \sigma'(a) = QaQ^{-1}.$$

Theorem (D.)

Twisting our bijection Ψ using σ' allows us to establish a bijection between:

- *Quantum adjacency operators $A \in \mathcal{B}(L^2(M))$;*
- *projections $e \in M \otimes M^{\text{op}}$ with $e = \sigma(e)$ and $(\sigma' \otimes \sigma')(e) = e$;*
- *self-adjoint M' -bimodules $S \subseteq \mathcal{B}(H)$ with $QSQ^{-1} = S$.*

So this is *more restrictive* than the tracial case.

Pullbacks

Let $\theta : M \rightarrow N$ be a normal CP map between von Neumann algebras $M \subseteq \mathcal{B}(H_M)$ and $N \subseteq \mathcal{B}(H_N)$. The Stinespring dilation takes a special form as θ is normal:

- there is a Hilbert space K and $U : H_N \rightarrow H_M \otimes K$;
- $\theta(x) = U^*(x \otimes 1)U$ for $x \in M \subseteq \mathcal{B}(H_M)$;
- there is a normal $*$ -homomorphism $\rho : N' \rightarrow H_M \otimes K$ with $Ux' = \rho(x')U$ for $x' \in N'$.

Given $S \subseteq \mathcal{B}(H_M)$ a Quantum (Graph/Relation) over M , define

$$\overleftarrow{S} = \text{weak}^*\text{-closure}\{U^*xU : x \in S \overline{\otimes} \mathcal{B}(K)\}.$$

Use of ρ shows that \overleftarrow{S} is a Quantum (Graph/Relation) over N , the “pullback”.

Pullbacks: Kraus forms

When M, N are finite-dimensional, $\theta : M \rightarrow N$ has a Kraus form

$$\theta(x) = \sum_{i=1}^n b_i^* x b_i.$$

(Notice I have swapped to considering UCP maps, not TPCP maps.)

Then [Weaver] for $S_1 \subseteq \mathcal{B}(H_M)$

$$\overleftarrow{S}_1 = \text{lin}\{b_i^* x b_j : x \in S_1\}.$$

Pushforwards

Given $S_2 \subseteq \mathcal{B}(H_N)$ a quantum relation over N , also

$$\overrightarrow{S_2} = \text{lin}\{b_i x b_j^* : x \in S_2\}$$

is a quantum relation over M , the “pushforward”.

Given classical graphs $G = (V_G, E_G)$ and $H = (V_H, E_H)$, a function $f : V_G \rightarrow V_H$ defines a $*$ -homomorphism (so certainly a UCP map)

$$\theta : C(V_H) \rightarrow C(V_G); \quad a \mapsto a \circ f \quad (a \in C(V_H)).$$

Let G induce $S_G \subseteq \mathcal{B}(\ell^2(V_G))$, that is,

$$S_G = \text{lin}\{e_{u,v} : (u,v) \in E_G\}$$

the span of matrix units supported on the edges. Then

$$\overrightarrow{S_G} = \text{lin}\{e_{f(u),f(v)} : (u,v) \in E_G\}$$

and so $\overrightarrow{S_G} \subseteq S_H$ exactly when f is a graph homomorphism.

Homomorphisms

[Stahkle] defines $\theta : M \rightarrow N$ to be a *homomorphism* between S_1 and S_2 when $\overrightarrow{S_2} \subseteq S_1$. [Weaver] calls this a *CP-morphism*.

Theorem (Stahkle)

Let G, H be (classical) graphs. Let $\theta : C(V_H) \rightarrow C(V_G)$ be a UCP map giving a homomorphism G to H (that is, with $\overrightarrow{S_G} \subseteq S_H$). Then there is some map $f : V_G \rightarrow V_H$ which is a (classical) homomorphism.

- In general θ need not be directly related to f .
- However, often we just care about the *existence* of a homomorphism.
- E.g. a k -colouring of G corresponds to some homomorphism $G \rightarrow K_k$, the complete graph. [I got confused in the talk, but this is correct!]

Further developments

The pushforward

$$\vec{S} = \text{lin}\{b_i x b_j^* : x \in S\}$$

doesn't make sense in the infinite-dimensional setting. [The definition exists, but it seems hard to prove anything.]

- What is a good notion of *homomorphism* in infinite dimensions?

Here we have worked exclusively with the operator bimodule picture of Quantum Graphs.

- Can we say something useful about homomorphisms and “adjacency matrices”?
- Already this seems problematic in the commutative case.

[Stop?]

Isomorphisms

Homomorphisms / CP-morphisms in this sense give a category.

Playing around with *multiplicative domains* for CP maps shows that the isomorphisms are exactly the $*$ -isomorphisms $\theta : M \rightarrow N$ which intertwine the Quantum Graphs.

- With $M \subseteq \mathcal{B}(L^2(M))$, any $*$ -automorphism $\theta : M \rightarrow M$ is implemented: there is a unitary $u \in \mathcal{B}(L^2(M))$ with $\theta(x) = u x u^*$.
- Then θ is an *automorphism* of the quantum graph S exactly when $u S u^* = S$.

What about an automorphism of the associated adjacency matrix A ?

- A acts on $L^2(M, \psi)$, say for some (tracial) ψ .
- It is hence natural to restrict to those θ which preserve ψ (automatic if ψ a trace).

Acting on $L^2(M)$

Have $\theta : M \rightarrow M$ a $*$ -automorphism.

Let $\Lambda : M \rightarrow L^2(M)$ be the GNS map. If θ preserves ψ then

$$\theta_0 : \Lambda(x) \mapsto \Lambda(\theta(x)) \quad (x \in M)$$

is an isometry (and so a unitary). (Indeed, θ_0 then implements θ .)
Then θ is an automorphism of our Quantum Graph if and only if

$$A\theta_0 = \theta_0 A \quad \text{on } L^2(M).$$

[Stop?]

Quantum Automorphisms

[We now shift gears...]

Let (A, Δ) be a compact quantum group. A *coaction* on M (still finite-dimensional) is a $*$ -homomorphism

$$\alpha : M \rightarrow M \otimes A; \quad (\alpha \otimes \text{id})\alpha = (\text{id} \otimes \Delta)\alpha,$$

and satisfying the density condition $\text{lin}\{(1 \otimes a)\alpha(x) : x \in M, a \in A\}$ dense in $M \otimes A$.

If we let M act on $L^2(M)$, then α has a *unitary implementation*, a unitary corepresentation $U \in \mathcal{B}(L^2(M)) \otimes A$ with

$$\alpha(x) = U(x \otimes 1)U^* \quad (x \in M).$$

We say that α coacts on the adjacency matrix A_G when

$$U(A_G \otimes 1) = (A_G \otimes 1)U.$$

Quantum Automorphisms of Operator Bimodules

$$U(A_G \otimes 1) = (A_G \otimes 1)U.$$

- Same as the single automorphism case, $uA_G = A_Gu$;
- Which corresponds to $uSu^* = S$

So might conjecture that we want $S \subseteq \mathcal{B}(L^2(M))$ and ask for $U(S \otimes 1)U^* = S \otimes 1$.

For various reasons, this doesn't work. For example, the “trivial quantum graph” is not preserved!

Instead, you need to twist by the *modular automorphism group*, or equivalently, look at a coaction of the *opposite quantum group*. Not clear to me exactly why this is...