

Around Compact Quantum Groups

Matthew Daws

23rd April 2009

What is a compact group?

Well, it's a compact topological space G with the structure of a group such that the group action is jointly continuous, and the inverse is continuous.

It's a unital commutative C^* -algebra A with a unital $*$ -homomorphism $\Delta : A \rightarrow A \otimes_{\min} A$ which is:

- ▶ Co-associative, $(\text{id} \otimes \Delta)\Delta = (\Delta \otimes \text{id})\Delta$
- ▶ “Cancellative”, that is, the sets

$$\{(a \otimes 1)\Delta(b) : a, b \in A\}, \quad \{(1 \otimes a)\Delta(b) : a, b \in A\},$$

have dense linear span in $A \otimes_{\min} A$.

What is a compact group?

Well, it's a compact topological space G with the structure of a group such that the group action is jointly continuous, and the inverse is continuous.

It's a unital commutative C^* -algebra A with a unital $*$ -homomorphism $\Delta : A \rightarrow A \otimes_{\min} A$ which is:

- ▶ Co-associative, $(\text{id} \otimes \Delta)\Delta = (\Delta \otimes \text{id})\Delta$
- ▶ “Cancellative”, that is, the sets

$$\{(a \otimes 1)\Delta(b) : a, b \in A\}, \quad \{(1 \otimes a)\Delta(b) : a, b \in A\},$$

have dense linear span in $A \otimes_{\min} A$.

What is a compact group?

Well, it's a compact topological space G with the structure of a group such that the group action is jointly continuous, and the inverse is continuous.

It's a unital commutative C^* -algebra A with a unital $*$ -homomorphism $\Delta : A \rightarrow A \otimes_{\min} A$ which is:

- ▶ Co-associative, $(\text{id} \otimes \Delta)\Delta = (\Delta \otimes \text{id})\Delta$
- ▶ “Cancellative”, that is, the sets

$$\{(a \otimes 1)\Delta(b) : a, b \in A\}, \quad \{(1 \otimes a)\Delta(b) : a, b \in A\},$$

have dense linear span in $A \otimes_{\min} A$.

What is a compact group?

Well, it's a compact topological space G with the structure of a group such that the group action is jointly continuous, and the inverse is continuous.

It's a unital commutative C^* -algebra A with a unital $*$ -homomorphism $\Delta : A \rightarrow A \otimes_{\min} A$ which is:

- ▶ Co-associative, $(\text{id} \otimes \Delta)\Delta = (\Delta \otimes \text{id})\Delta$
- ▶ “Cancellative”, that is, the sets

$$\{(a \otimes 1)\Delta(b) : a, b \in A\}, \quad \{(1 \otimes a)\Delta(b) : a, b \in A\},$$

have dense linear span in $A \otimes_{\min} A$.

Equivalence, easy direction

If G is a compact group, set

$$A = C(G) = \{\text{continuous functions } G \rightarrow \mathbb{C}\},$$

identify $A \otimes_{\min} A = C(G \times G)$, define

$$\Delta(f) \in C(G \times G), \quad \Delta(f) : (s, t) \mapsto f(st) \quad (f \in C(G), s, t \in G).$$

Finally observe that

$$(a \otimes 1)\Delta(b) : (s, t) \mapsto a(s)b(st),$$

will separate the points of $G \times G$ (by varying a and b) so by Stone-Weierstrass,

$$\text{lin}\{(a \otimes 1)\Delta(b) : a, b \in A\}$$

is a dense subalgebra of $C(G \times G)$.

Equivalence, easy direction

If G is a compact group, set

$$A = C(G) = \{\text{continuous functions } G \rightarrow \mathbb{C}\},$$

identify $A \otimes_{\min} A = C(G \times G)$, define

$$\Delta(f) \in C(G \times G), \quad \Delta(f) : (s, t) \mapsto f(st) \quad (f \in C(G), s, t \in G).$$

Finally observe that

$$(a \otimes 1)\Delta(b) : (s, t) \mapsto a(s)b(st),$$

will separate the points of $G \times G$ (by varying a and b) so by Stone-Weierstrass,

$$\text{lin}\{(a \otimes 1)\Delta(b) : a, b \in A\}$$

is a dense subalgebra of $C(G \times G)$.

Equivalence, easy direction

If G is a compact group, set

$$A = C(G) = \{\text{continuous functions } G \rightarrow \mathbb{C}\},$$

identify $A \otimes_{\min} A = C(G \times G)$, define

$$\Delta(f) \in C(G \times G), \quad \Delta(f) : (s, t) \mapsto f(st) \quad (f \in C(G), s, t \in G).$$

Finally observe that

$$(a \otimes 1)\Delta(b) : (s, t) \mapsto a(s)b(st),$$

will separate the points of $G \times G$ (by varying a and b) so by Stone-Weierstrass,

$$\text{lin}\{(a \otimes 1)\Delta(b) : a, b \in A\}$$

is a dense subalgebra of $C(G \times G)$.

Equivalence, easy direction

If G is a compact group, set

$$A = C(G) = \{\text{continuous functions } G \rightarrow \mathbb{C}\},$$

identify $A \otimes_{\min} A = C(G \times G)$, define

$$\Delta(f) \in C(G \times G), \quad \Delta(f) : (s, t) \mapsto f(st) \quad (f \in C(G), s, t \in G).$$

Finally observe that

$$(a \otimes 1)\Delta(b) : (s, t) \mapsto a(s)b(st),$$

will separate the points of $G \times G$ (by varying a and b) so by Stone-Weierstrass,

$$\text{lin}\{(a \otimes 1)\Delta(b) : a, b \in A\}$$

is a dense subalgebra of $C(G \times G)$.

Equivalence, hard direction

Gelfand-Naimark tells us that a unital commutative C^* -algebra A has the form $C(X)$ for some compact space X . So again $A \otimes_{\min} A = C(X \times X)$. Then $\Delta : C(X) \rightarrow C(X \times X)$ a unital $*$ -homomorphism induces a continuous map $\theta : X \times X \rightarrow X$ such that

$$f(\theta(s, t)) = \Delta(f)(s, t) \quad (s, t \in X, f \in C(X)).$$

The category of unital commutative C^* -algebras and unital $*$ -homomorphisms is dual to the category of compact spaces and continuous maps.

Δ co-associative implies that θ is associative, so X is a compact semigroup.

The cancellation rules for Δ imply that X is cancellative, that is

$$st = rt \implies s = r, \quad ts = tr \implies s = r.$$

Exercise: A compact semigroup with cancellation is a compact group.

Equivalence, hard direction

Gelfand-Naimark tells us that a unital commutative C^* -algebra A has the form $C(X)$ for some compact space X . So again $A \otimes_{\min} A = C(X \times X)$. Then $\Delta : C(X) \rightarrow C(X \times X)$ a unital $*$ -homomorphism induces a continuous map $\theta : X \times X \rightarrow X$ such that

$$f(\theta(s, t)) = \Delta(f)(s, t) \quad (s, t \in X, f \in C(X)).$$

The category of unital commutative C^* -algebras and unital $*$ -homomorphisms is dual to the category of compact spaces and continuous maps.

Δ co-associative implies that θ is associative, so X is a compact semigroup.

The cancellation rules for Δ imply that X is cancellative, that is

$$st = rt \implies s = r, \quad ts = tr \implies s = r.$$

Exercise: A compact semigroup with cancellation is a compact group.

Equivalence, hard direction

Gelfand-Naimark tells us that a unital commutative C^* -algebra A has the form $C(X)$ for some compact space X . So again $A \otimes_{\min} A = C(X \times X)$. Then $\Delta : C(X) \rightarrow C(X \times X)$ a unital $*$ -homomorphism induces a continuous map $\theta : X \times X \rightarrow X$ such that

$$f(\theta(s, t)) = \Delta(f)(s, t) \quad (s, t \in X, f \in C(X)).$$

The category of unital commutative C^* -algebras and unital $*$ -homomorphisms is dual to the category of compact spaces and continuous maps.

Δ co-associative implies that θ is associative, so X is a compact semigroup.

The cancellation rules for Δ imply that X is cancellative, that is

$$st = rt \implies s = r, \quad ts = tr \implies s = r.$$

Exercise: A compact semigroup with cancellation is a compact group.

Equivalence, hard direction

Gelfand-Naimark tells us that a unital commutative C^* -algebra A has the form $C(X)$ for some compact space X . So again $A \otimes_{\min} A = C(X \times X)$. Then $\Delta : C(X) \rightarrow C(X \times X)$ a unital $*$ -homomorphism induces a continuous map $\theta : X \times X \rightarrow X$ such that

$$f(\theta(s, t)) = \Delta(f)(s, t) \quad (s, t \in X, f \in C(X)).$$

The category of unital commutative C^* -algebras and unital $*$ -homomorphisms is dual to the category of compact spaces and continuous maps.

Δ co-associative implies that θ is associative, so X is a compact semigroup.

The cancellation rules for Δ imply that X is cancellative, that is

$$st = rt \implies s = r, \quad ts = tr \implies s = r.$$

Exercise: A compact semigroup with cancellation is a compact group.

Equivalence, hard direction

Gelfand-Naimark tells us that a unital commutative C^* -algebra A has the form $C(X)$ for some compact space X . So again $A \otimes_{\min} A = C(X \times X)$. Then $\Delta : C(X) \rightarrow C(X \times X)$ a unital $*$ -homomorphism induces a continuous map $\theta : X \times X \rightarrow X$ such that

$$f(\theta(s, t)) = \Delta(f)(s, t) \quad (s, t \in X, f \in C(X)).$$

The category of unital commutative C^* -algebras and unital $*$ -homomorphisms is dual to the category of compact spaces and continuous maps.

Δ co-associative implies that θ is associative, so X is a compact semigroup.

The cancellation rules for Δ imply that X is cancellative, that is

$$st = rt \implies s = r, \quad ts = tr \implies s = r.$$

Exercise: A compact semigroup with cancellation is a compact group.

Equivalence, hard direction

Gelfand-Naimark tells us that a unital commutative C^* -algebra A has the form $C(X)$ for some compact space X . So again $A \otimes_{\min} A = C(X \times X)$. Then $\Delta : C(X) \rightarrow C(X \times X)$ a unital $*$ -homomorphism induces a continuous map $\theta : X \times X \rightarrow X$ such that

$$f(\theta(s, t)) = \Delta(f)(s, t) \quad (s, t \in X, f \in C(X)).$$

The category of unital commutative C^* -algebras and unital $*$ -homomorphisms is dual to the category of compact spaces and continuous maps.

Δ co-associative implies that θ is associative, so X is a compact semigroup.

The cancellation rules for Δ imply that X is cancellative, that is

$$st = rt \implies s = r, \quad ts = tr \implies s = r.$$

Exercise: A compact semigroup with cancellation is a compact group.

Compact quantum groups

Simply remove the word “commutative”!

For example, let Γ be a discrete group, and let Γ act on $\ell^2(\Gamma)$ by left translation:

$$\lambda(s)f : t \mapsto f(s^{-1}t) \quad (s, t \in \Gamma, f \in \ell^2(\Gamma)).$$

Let $C_r^*(\Gamma)$ be the (reduced) group C^* -algebra: that is, the norm closed algebra, acting on $\ell^2(\Gamma)$, generated by $\lambda(\Gamma)$. So $C_r^*(\Gamma)$ is commutative if and only if Γ is.

There is a $*$ -homomorphism

$$\begin{aligned} \Delta : C_r^*(\Gamma) &\rightarrow C_r^*(\Gamma) \otimes_{\min} C_r^*(\Gamma) = C_r^*(\Gamma \times \Gamma), \\ \Delta : \lambda(s) &\mapsto \lambda(s) \otimes \lambda(s) = \lambda(s, s) \quad (s \in \Gamma). \end{aligned}$$

Compact quantum groups

Simply remove the word “commutative”!

For example, let Γ be a discrete group, and let Γ act on $\ell^2(\Gamma)$ by left translation:

$$\lambda(s)f : t \mapsto f(s^{-1}t) \quad (s, t \in \Gamma, f \in \ell^2(\Gamma)).$$

Let $C_r^*(\Gamma)$ be the (reduced) group C^* -algebra: that is, the norm closed algebra, acting on $\ell^2(\Gamma)$, generated by $\lambda(\Gamma)$. So $C_r^*(\Gamma)$ is commutative if and only if Γ is.

There is a $*$ -homomorphism

$$\begin{aligned} \Delta : C_r^*(\Gamma) &\rightarrow C_r^*(\Gamma) \otimes_{\min} C_r^*(\Gamma) = C_r^*(\Gamma \times \Gamma), \\ \Delta : \lambda(s) &\mapsto \lambda(s) \otimes \lambda(s) = \lambda(s, s) \quad (s \in \Gamma). \end{aligned}$$

Compact quantum groups

Simply remove the word “commutative”!

For example, let Γ be a discrete group, and let Γ act on $\ell^2(\Gamma)$ by left translation:

$$\lambda(s)f : t \mapsto f(s^{-1}t) \quad (s, t \in \Gamma, f \in \ell^2(\Gamma)).$$

Let $C_r^*(\Gamma)$ be the (reduced) group C^* -algebra: that is, the norm closed algebra, acting on $\ell^2(\Gamma)$, generated by $\lambda(\Gamma)$. So $C_r^*(\Gamma)$ is commutative if and only if Γ is.

There is a $*$ -homomorphism

$$\begin{aligned} \Delta : C_r^*(\Gamma) &\rightarrow C_r^*(\Gamma) \otimes_{\min} C_r^*(\Gamma) = C_r^*(\Gamma \times \Gamma), \\ \Delta : \lambda(s) &\mapsto \lambda(s) \otimes \lambda(s) = \lambda(s, s) \quad (s \in \Gamma). \end{aligned}$$

Compact quantum groups

Simply remove the word “commutative”!

For example, let Γ be a discrete group, and let Γ act on $\ell^2(\Gamma)$ by left translation:

$$\lambda(s)f : t \mapsto f(s^{-1}t) \quad (s, t \in \Gamma, f \in \ell^2(\Gamma)).$$

Let $C_r^*(\Gamma)$ be the (reduced) group C^* -algebra: that is, the norm closed algebra, acting on $\ell^2(\Gamma)$, generated by $\lambda(\Gamma)$. So $C_r^*(\Gamma)$ is commutative if and only if Γ is.

There is a $*$ -homomorphism

$$\begin{aligned} \Delta : C_r^*(\Gamma) &\rightarrow C_r^*(\Gamma) \otimes_{\min} C_r^*(\Gamma) = C_r^*(\Gamma \times \Gamma), \\ \Delta : \lambda(s) &\mapsto \lambda(s) \otimes \lambda(s) = \lambda(s, s) \quad (s \in \Gamma). \end{aligned}$$

Compact quantum groups

Simply remove the word “commutative”!

For example, let Γ be a discrete group, and let Γ act on $\ell^2(\Gamma)$ by left translation:

$$\lambda(s)f : t \mapsto f(s^{-1}t) \quad (s, t \in \Gamma, f \in \ell^2(\Gamma)).$$

Let $C_r^*(\Gamma)$ be the (reduced) group C^* -algebra: that is, the norm closed algebra, acting on $\ell^2(\Gamma)$, generated by $\lambda(\Gamma)$. So $C_r^*(\Gamma)$ is commutative if and only if Γ is.

There is a $*$ -homomorphism

$$\begin{aligned} \Delta : C_r^*(\Gamma) &\rightarrow C_r^*(\Gamma) \otimes_{\min} C_r^*(\Gamma) = C_r^*(\Gamma \times \Gamma), \\ \Delta : \lambda(s) &\mapsto \lambda(s) \otimes \lambda(s) = \lambda(s, s) \quad (s \in \Gamma). \end{aligned}$$

Compact or Discrete?

Hang on: we're saying that for *discrete* Γ , we have that $C_r^*(\Gamma)$ is a *compact* quantum group?

If Γ were abelian, then the fourier transform tells us that

$$C_r^*(\Gamma) \cong C(\hat{\Gamma}),$$

where $\hat{\Gamma}$ is the Pontryagin dual of Γ . As Γ is discrete, $\hat{\Gamma}$ is compact.

As $C(G)$ is our “commutative” base algebra, this weird terminology is forced upon us.

Compact or Discrete?

Hang on: we're saying that for *discrete* Γ , we have that $C_r^*(\Gamma)$ is a *compact* quantum group?

If Γ were abelian, then the fourier transform tells us that

$$C_r^*(\Gamma) \cong C(\hat{\Gamma}),$$

where $\hat{\Gamma}$ is the Pontryagin dual of Γ . As Γ is discrete, $\hat{\Gamma}$ is compact.

As $C(G)$ is our “commutative” base algebra, this weird terminology is forced upon us.

Compact or Discrete?

Hang on: we're saying that for *discrete* Γ , we have that $C_r^*(\Gamma)$ is a *compact* quantum group?

If Γ were abelian, then the fourier transform tells us that

$$C_r^*(\Gamma) \cong C(\hat{\Gamma}),$$

where $\hat{\Gamma}$ is the Pontryagin dual of Γ . As Γ is discrete, $\hat{\Gamma}$ is compact.

As $C(G)$ is our “commutative” base algebra, this weird terminology is forced upon us.

Compact or Discrete?

Hang on: we're saying that for *discrete* Γ , we have that $C_r^*(\Gamma)$ is a *compact* quantum group?

If Γ were abelian, then the fourier transform tells us that

$$C_r^*(\Gamma) \cong C(\hat{\Gamma}),$$

where $\hat{\Gamma}$ is the Pontryagin dual of Γ . As Γ is discrete, $\hat{\Gamma}$ is compact.

As $C(G)$ is our “commutative” base algebra, this weird terminology is forced upon us.

Twisted $SU(2)$

From Woronowicz in the C^* -setting, but independently discovered by Soibelman and Vaksman

$C(SU(2))$ is the commutative C^* -algebra generated by a, b with

$$a^*a + b^*b = 1.$$

$$aa^* + bb^* = 1, \quad b^*b = bb^*, \quad ab = ba, \quad ab^* = b^*a.$$

We introduce a real parameter $\mu \in [-1, 1] \setminus \{0\}$, and let $C(SU_\mu(2))$ be the (non-commutative) C^* -algebra generated by a, b with

$$\begin{aligned} a^*a + b^*b &= 1, & aa^* + \mu^2 bb^* &= 1, \\ b^*b &= bb^*, & ab &= \mu ba, & ab^* &= \mu b^*a. \end{aligned}$$

There exists a coproduct Δ with

$$\Delta(a) = a \otimes a - \mu b^* \otimes b, \quad \Delta(b) = b \otimes a + a^* \otimes b.$$

Twisted $SU(2)$

From Woronowicz in the C^* -setting, but independently discovered by Soibelman and Vaksman

$C(SU(2))$ is the commutative C^* -algebra generated by a, b with

$$a^*a + b^*b = 1, \\ aa^* + bb^* = 1, \quad b^*b = bb^*, \quad ab = ba, \quad ab^* = b^*a.$$

We introduce a real parameter $\mu \in [-1, 1] \setminus \{0\}$, and let $C(SU_\mu(2))$ be the (non-commutative) C^* -algebra generated by a, b with

$$a^*a + b^*b = 1, \quad aa^* + \mu^2 bb^* = 1, \\ b^*b = bb^*, \quad ab = \mu ba, \quad ab^* = \mu b^*a.$$

There exists a coproduct Δ with

$$\Delta(a) = a \otimes a - \mu b^* \otimes b, \quad \Delta(b) = b \otimes a + a^* \otimes b.$$

Twisted $SU(2)$

From Woronowicz in the C^* -setting, but independently discovered by Soibelman and Vaksman

$C(SU(2))$ is the commutative C^* -algebra generated by a, b with

$$a^*a + b^*b = 1, \\ aa^* + bb^* = 1, \quad b^*b = bb^*, \quad ab = ba, \quad ab^* = b^*a.$$

We introduce a real parameter $\mu \in [-1, 1] \setminus \{0\}$, and let $C(SU_\mu(2))$ be the (non-commutative) C^* -algebra generated by a, b with

$$a^*a + b^*b = 1, \quad aa^* + \mu^2 bb^* = 1, \\ b^*b = bb^*, \quad ab = \mu ba, \quad ab^* = \mu b^*a.$$

There exists a coproduct Δ with

$$\Delta(a) = a \otimes a - \mu b^* \otimes b, \quad \Delta(b) = b \otimes a + a^* \otimes b.$$

Twisted $SU(2)$

From Woronowicz in the C^* -setting, but independently discovered by Soibelman and Vaksman

$C(SU(2))$ is the commutative C^* -algebra generated by a, b with

$$\begin{aligned} a^*a + b^*b &= 1, \\ aa^* + bb^* &= 1, \quad b^*b = bb^*, \quad ab = ba, \quad ab^* = b^*a. \end{aligned}$$

We introduce a real parameter $\mu \in [-1, 1] \setminus \{0\}$, and let $C(SU_\mu(2))$ be the (non-commutative) C^* -algebra generated by a, b with

$$\begin{aligned} a^*a + b^*b &= 1, \quad aa^* + \mu^2bb^* = 1, \\ b^*b &= bb^*, \quad ab = \mu ba, \quad ab^* = \mu b^*a. \end{aligned}$$

There exists a coproduct Δ with

$$\Delta(a) = a \otimes a - \mu b^* \otimes b, \quad \Delta(b) = b \otimes a + a^* \otimes b.$$

Why?

(Topological) Quantum groups grew out of:

- ▶ How do we extend the notion of the Fourier transform, or more specifically, the Pontryagin Duality, to non-abelian groups? Ideally, we'd like a self-dual category into which all proper groups fit (so Tannaka-Krein duality doesn't quite fit the bill). This lead to *Kac algebras*.
- ▶ But $SU_\mu(2)$ does not fit into this framework! Indeed, we have rather few examples of Kac algebras.

We now have a simple set of axioms for objects which are called “locally compact quantum groups”, and which encompass all known examples.

Why?

(Topological) Quantum groups grew out of:

- ▶ How do we extend the notion of the Fourier transform, or more specifically, the Pontryagin Duality, to non-abelian groups? Ideally, we'd like a self-dual category into which all proper groups fit (so Tannaka-Krein duality doesn't quite fit the bill). This led to *Kac algebras*.
- ▶ But $SU_\mu(2)$ does not fit into this framework! Indeed, we have rather few examples of Kac algebras.

We now have a simple set of axioms for objects which are called “locally compact quantum groups”, and which encompass all known examples.

Why?

(Topological) Quantum groups grew out of:

- ▶ How do we extend the notion of the Fourier transform, or more specifically, the Pontryagin Duality, to non-abelian groups? Ideally, we'd like a self-dual category into which all proper groups fit (so Tannaka-Krein duality doesn't quite fit the bill). This lead to *Kac algebras*.
- ▶ But $SU_\mu(2)$ does not fit into this framework! Indeed, we have rather few examples of Kac algebras.

We now have a simple set of axioms for objects which are called “locally compact quantum groups”, and which encompass all known examples.

Why?

(Topological) Quantum groups grew out of:

- ▶ How do we extend the notion of the Fourier transform, or more specifically, the Pontryagin Duality, to non-abelian groups? Ideally, we'd like a self-dual category into which all proper groups fit (so Tannaka-Krein duality doesn't quite fit the bill). This lead to *Kac algebras*.
- ▶ But $SU_\mu(2)$ does not fit into this framework! Indeed, we have rather few examples of Kac algebras.

We now have a simple set of axioms for objects which are called “locally compact quantum groups”, and which encompass all known examples.

Why?

(Topological) Quantum groups grew out of:

- ▶ How do we extend the notion of the Fourier transform, or more specifically, the Pontryagin Duality, to non-abelian groups? Ideally, we'd like a self-dual category into which all proper groups fit (so Tannaka-Krein duality doesn't quite fit the bill). This lead to *Kac algebras*.
- ▶ But $SU_{\mu}(2)$ does not fit into this framework! Indeed, we have rather few examples of Kac algebras.

We now have a simple set of axioms for objects which are called “locally compact quantum groups”, and which encompass all known examples.

Why?

(Topological) Quantum groups grew out of:

- ▶ How do we extend the notion of the Fourier transform, or more specifically, the Pontryagin Duality, to non-abelian groups? Ideally, we'd like a self-dual category into which all proper groups fit (so Tannaka-Krein duality doesn't quite fit the bill). This lead to *Kac algebras*.
- ▶ But $SU_{\mu}(2)$ does not fit into this framework! Indeed, we have rather few examples of Kac algebras.

We now have a simple set of axioms for objects which are called “locally compact quantum groups”, and which encompass all known examples.

Why?

(Topological) Quantum groups grew out of:

- ▶ How do we extend the notion of the Fourier transform, or more specifically, the Pontryagin Duality, to non-abelian groups? Ideally, we'd like a self-dual category into which all proper groups fit (so Tannaka-Krein duality doesn't quite fit the bill). This lead to *Kac algebras*.
- ▶ But $SU_{\mu}(2)$ does not fit into this framework! Indeed, we have rather few examples of Kac algebras.

We now have a simple set of axioms for objects which are called “locally compact quantum groups”, and which encompass all known examples.

Haar state

A compact group admits a unique Haar measure: a probability measure which is invariant under the group action. In our language, this corresponds to a *state* φ on $C(G)$ with

$$(\varphi \otimes \text{id})\Delta(a) = (\text{id} \otimes \varphi)\Delta(a) = \varphi(a) \quad (a \in C(G)).$$

Woronowicz and van Daele showed that such a state always exists on a compact quantum group (A, Δ)

Applying the GNS construction gives a Hilbert space H and a $*$ -representation of A on H . This is the analogue of $C(G)$ acting on $L^2(G)$ by pointwise multiplication.

Haar state

A compact group admits a unique Haar measure: a probability measure which is invariant under the group action. In our language, this corresponds to a *state* φ on $C(G)$ with

$$(\varphi \otimes \text{id})\Delta(a) = (\text{id} \otimes \varphi)\Delta(a) = \varphi(a) \quad (a \in C(G)).$$

Woronowicz and van Daele showed that such a state always exists on a compact quantum group (A, Δ) . Applying the GNS construction gives a Hilbert space H and a $*$ -representation of A on H . This is the analogue of $C(G)$ acting on $L^2(G)$ by pointwise multiplication.

Haar state

A compact group admits a unique Haar measure: a probability measure which is invariant under the group action. In our language, this corresponds to a *state* φ on $C(G)$ with

$$(\varphi \otimes \text{id})\Delta(a) = (\text{id} \otimes \varphi)\Delta(a) = \varphi(a) \quad (a \in C(G)).$$

Woronowicz and van Daele showed that such a state always exists on a compact quantum group (A, Δ) . Applying the GNS construction gives a Hilbert space H and a $*$ -representation of A on H . This is the analogue of $C(G)$ acting on $L^2(G)$ by pointwise multiplication.

Haar state

A compact group admits a unique Haar measure: a probability measure which is invariant under the group action. In our language, this corresponds to a *state* φ on $C(G)$ with

$$(\varphi \otimes \text{id})\Delta(a) = (\text{id} \otimes \varphi)\Delta(a) = \varphi(a) \quad (a \in C(G)).$$

Woronowicz and van Daele showed that such a state always exists on a compact quantum group (A, Δ)

Applying the GNS construction gives a Hilbert space H and a $*$ -representation of A on H . This is the analogue of $C(G)$ acting on $L^2(G)$ by pointwise multiplication.

Haar state

A compact group admits a unique Haar measure: a probability measure which is invariant under the group action. In our language, this corresponds to a *state* φ on $C(G)$ with

$$(\varphi \otimes \text{id})\Delta(a) = (\text{id} \otimes \varphi)\Delta(a) = \varphi(a) \quad (a \in C(G)).$$

Woronowicz and van Daele showed that such a state always exists on a compact quantum group (A, Δ)

Applying the GNS construction gives a Hilbert space H and a $*$ -representation of A on H . This is the analogue of $C(G)$ acting on $L^2(G)$ by pointwise multiplication.

Haar state

A compact group admits a unique Haar measure: a probability measure which is invariant under the group action. In our language, this corresponds to a *state* φ on $C(G)$ with

$$(\varphi \otimes \text{id})\Delta(a) = (\text{id} \otimes \varphi)\Delta(a) = \varphi(a) \quad (a \in C(G)).$$

Woronowicz and van Daele showed that such a state always exists on a compact quantum group (A, Δ)

Applying the GNS construction gives a Hilbert space H and a $*$ -representation of A on H . This is the analogue of $C(G)$ acting on $L^2(G)$ by pointwise multiplication.

Corepresentation theory

A (finite-dimensional) corepresentation of (A, Δ) is a matrix $u \in \mathbb{M}_n(A)$ with

$$\Delta(u_{ij}) = \sum_{k=1}^n u_{ik} \otimes u_{kj}.$$

- ▶ All irreducible corepresentations of (A, Δ) are finite-dimensional.
- ▶ Using the Haar state, it's possible to show that any finite-dimensional corepresentation u is *equivalent* to a unitary corepresentation: $u^*u = uu^* = I_n$.
- ▶ There is an (infinite-dimensional) corepresentation of (A, Δ) on H ; all finite-dimensional corepresentations are sub-corepresentations of this. So we have a generalised Peter-Weil theory.
- ▶ There is a character theory.

Corepresentation theory

A (finite-dimensional) corepresentation of (A, Δ) is a matrix $u \in \mathbb{M}_n(A)$ with

$$\Delta(u_{ij}) = \sum_{k=1}^n u_{ik} \otimes u_{kj}.$$

- ▶ All irreducible corepresentations of (A, Δ) are finite-dimensional.
- ▶ Using the Haar state, it's possible to show that any finite-dimensional corepresentation u is *equivalent* to a unitary corepresentation: $u^*u = uu^* = I_n$.
- ▶ There is an (infinite-dimensional) corepresentation of (A, Δ) on H ; all finite-dimensional corepresentations are sub-corepresentations of this. So we have a generalised Peter-Weil theory.
- ▶ There is a character theory.

Corepresentation theory

A (finite-dimensional) corepresentation of (A, Δ) is a matrix $u \in \mathbb{M}_n(A)$ with

$$\Delta(u_{ij}) = \sum_{k=1}^n u_{ik} \otimes u_{kj}.$$

- ▶ All irreducible corepresentations of (A, Δ) are finite-dimensional.
- ▶ Using the Haar state, it's possible to show that any finite-dimensional corepresentation u is *equivalent* to a unitary corepresentation: $u^*u = uu^* = I_n$.
- ▶ There is an (infinite-dimensional) corepresentation of (A, Δ) on H ; all finite-dimensional corepresentations are sub-corepresentations of this. So we have a generalised Peter-Weil theory.
- ▶ There is a character theory.

Corepresentation theory

A (finite-dimensional) corepresentation of (A, Δ) is a matrix $u \in \mathbb{M}_n(A)$ with

$$\Delta(u_{ij}) = \sum_{k=1}^n u_{ik} \otimes u_{kj}.$$

- ▶ All irreducible corepresentations of (A, Δ) are finite-dimensional.
- ▶ Using the Haar state, it's possible to show that any finite-dimensional corepresentation u is *equivalent* to a unitary corepresentation: $u^*u = uu^* = I_n$.
- ▶ There is an (infinite-dimensional) corepresentation of (A, Δ) on H ; all finite-dimensional corepresentations are sub-corepresentations of this. So we have a generalised Peter-Weil theory.
- ▶ There is a character theory.

Corepresentation theory

A (finite-dimensional) corepresentation of (A, Δ) is a matrix $u \in \mathbb{M}_n(A)$ with

$$\Delta(u_{ij}) = \sum_{k=1}^n u_{ik} \otimes u_{kj}.$$

- ▶ All irreducible corepresentations of (A, Δ) are finite-dimensional.
- ▶ Using the Haar state, it's possible to show that any finite-dimensional corepresentation u is *equivalent* to a unitary corepresentation: $u^*u = uu^* = I_n$.
- ▶ There is an (infinite-dimensional) corepresentation of (A, Δ) on H ; all finite-dimensional corepresentations are sub-corepresentations of this. So we have a generalised Peter-Weil theory.
- ▶ There is a character theory.

Corepresentation theory

A (finite-dimensional) corepresentation of (A, Δ) is a matrix $u \in \mathbb{M}_n(A)$ with

$$\Delta(u_{ij}) = \sum_{k=1}^n u_{ik} \otimes u_{kj}.$$

- ▶ All irreducible corepresentations of (A, Δ) are finite-dimensional.
- ▶ Using the Haar state, it's possible to show that any finite-dimensional corepresentation u is *equivalent* to a unitary corepresentation: $u^*u = uu^* = I_n$.
- ▶ There is an (infinite-dimensional) corepresentation of (A, Δ) on H ; all finite-dimensional corepresentations are sub-corepresentations of this. So we have a generalised Peter-Weil theory.
- ▶ There is a character theory.

Corepresentation theory

A (finite-dimensional) corepresentation of (A, Δ) is a matrix $u \in \mathbb{M}_n(A)$ with

$$\Delta(u_{ij}) = \sum_{k=1}^n u_{ik} \otimes u_{kj}.$$

- ▶ All irreducible corepresentations of (A, Δ) are finite-dimensional.
- ▶ Using the Haar state, it's possible to show that any finite-dimensional corepresentation u is *equivalent* to a unitary corepresentation: $u^*u = uu^* = I_n$.
- ▶ There is an (infinite-dimensional) corepresentation of (A, Δ) on H ; all finite-dimensional corepresentations are sub-corepresentations of this. So we have a generalised Peter-Weil theory.
- ▶ There is a character theory.

Links with algebra

Let \mathcal{A} be the collection of matrix entries of all irreducible corepresentations of (A, Δ) .

- ▶ Then \mathcal{A} is a $*$ -algebra.
- ▶ \mathcal{A} is norm-dense in A .
- ▶ Δ restricts to give a map $\mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$ (algebraic tensor product).
- ▶ We can turn (\mathcal{A}, Δ) into a Hopf $*$ -algebra: there exists an antipode and counit.
- ▶ But, in general, these are *unbounded*, and so don't make sense on A .
- ▶ (\mathcal{A}, Δ) is unique.

Links with algebra

Let \mathcal{A} be the collection of matrix entries of all irreducible corepresentations of (A, Δ) .

- ▶ Then \mathcal{A} is a $*$ -algebra.
- ▶ \mathcal{A} is norm-dense in A .
- ▶ Δ restricts to give a map $\mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$ (algebraic tensor product).
- ▶ We can turn (\mathcal{A}, Δ) into a Hopf $*$ -algebra: there exists an antipode and counit.
- ▶ But, in general, these are *unbounded*, and so don't make sense on A .
- ▶ (\mathcal{A}, Δ) is unique.

Links with algebra

Let \mathcal{A} be the collection of matrix entries of all irreducible corepresentations of (A, Δ) .

- ▶ Then \mathcal{A} is a $*$ -algebra.
- ▶ \mathcal{A} is norm-dense in A .
- ▶ Δ restricts to give a map $\mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$ (algebraic tensor product).
- ▶ We can turn (\mathcal{A}, Δ) into a Hopf $*$ -algebra: there exists an antipode and counit.
- ▶ But, in general, these are *unbounded*, and so don't make sense on A .
- ▶ (\mathcal{A}, Δ) is unique.

Links with algebra

Let \mathcal{A} be the collection of matrix entries of all irreducible corepresentations of (A, Δ) .

- ▶ Then \mathcal{A} is a $*$ -algebra.
- ▶ \mathcal{A} is norm-dense in A .
- ▶ Δ restricts to give a map $\mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$ (algebraic tensor product).
- ▶ We can turn (\mathcal{A}, Δ) into a Hopf $*$ -algebra: there exists an antipode and counit.
- ▶ But, in general, these are *unbounded*, and so don't make sense on A .
- ▶ (\mathcal{A}, Δ) is unique.

Links with algebra

Let \mathcal{A} be the collection of matrix entries of all irreducible corepresentations of (A, Δ) .

- ▶ Then \mathcal{A} is a $*$ -algebra.
- ▶ \mathcal{A} is norm-dense in A .
- ▶ Δ restricts to give a map $\mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$ (algebraic tensor product).
- ▶ We can turn (\mathcal{A}, Δ) into a Hopf $*$ -algebra: there exists an antipode and counit.
- ▶ But, in general, these are *unbounded*, and so don't make sense on A .
- ▶ (\mathcal{A}, Δ) is unique.

Links with algebra

Let \mathcal{A} be the collection of matrix entries of all irreducible corepresentations of (A, Δ) .

- ▶ Then \mathcal{A} is a $*$ -algebra.
- ▶ \mathcal{A} is norm-dense in A .
- ▶ Δ restricts to give a map $\mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$ (algebraic tensor product).
- ▶ We can turn (\mathcal{A}, Δ) into a Hopf $*$ -algebra: there exists an antipode and counit.
- ▶ But, in general, these are *unbounded*, and so don't make sense on A .
- ▶ (\mathcal{A}, Δ) is unique.

Links with algebra

Let \mathcal{A} be the collection of matrix entries of all irreducible corepresentations of (A, Δ) .

- ▶ Then \mathcal{A} is a $*$ -algebra.
- ▶ \mathcal{A} is norm-dense in A .
- ▶ Δ restricts to give a map $\mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$ (algebraic tensor product).
- ▶ We can turn (\mathcal{A}, Δ) into a Hopf $*$ -algebra: there exists an antipode and counit.
- ▶ But, in general, these are *unbounded*, and so don't make sense on A .
- ▶ (\mathcal{A}, Δ) is unique.

Links with algebra

Let \mathcal{A} be the collection of matrix entries of all irreducible corepresentations of (A, Δ) .

- ▶ Then \mathcal{A} is a $*$ -algebra.
- ▶ \mathcal{A} is norm-dense in A .
- ▶ Δ restricts to give a map $\mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$ (algebraic tensor product).
- ▶ We can turn (\mathcal{A}, Δ) into a Hopf $*$ -algebra: there exists an antipode and counit.
- ▶ But, in general, these are *unbounded*, and so don't make sense on A .
- ▶ (\mathcal{A}, Δ) is unique.

More algebra

We can go in reverse! Dijkhuizen and Koornwinder showed that if (\mathcal{A}, Δ) is a Hopf $*$ -algebra which is spanned by the matrix entries of its finite-dimensional unitary corepresentations, then there is a compact quantum group (A, Δ) such that \mathcal{A} is given by A .

Drinfeld's approach to quantum groups starts with a Lie group G , the Lie algebra \mathfrak{g} and the enveloping algebra $U(\mathfrak{g})$ which is naturally a Hopf algebra. It is this Hopf algebra which is deformed to get a "quantum group". If we start with $SU(2)$, and we now take the $*$ -representations of this deformed enveloping algebra, we naturally get a Hopf $*$ -algebra which is isomorphic to the Hopf $*$ -algebra associated to $C(SU_\mu(G))$.

So we are really studying the "dual" world to what is often understood by the term "quantum group".

More algebra

We can go in reverse! Dijkhuizen and Koornwinder showed that if (\mathcal{A}, Δ) is a Hopf $*$ -algebra which is spanned by the matrix entries of its finite-dimensional unitary corepresentations, then there is a compact quantum group (A, Δ) such that \mathcal{A} is given by A .

Drinfeld's approach to quantum groups starts with a Lie group G , the Lie algebra \mathfrak{g} and the enveloping algebra $U(\mathfrak{g})$ which is naturally a Hopf algebra. It is this Hopf algebra which is deformed to get a "quantum group". If we start with $SU(2)$, and we now take the $*$ -representations of this deformed enveloping algebra, we naturally get a Hopf $*$ -algebra which is isomorphic to the Hopf $*$ -algebra associated to $C(SU_\mu(G))$.

So we are really studying the "dual" world to what is often understood by the term "quantum group".

More algebra

We can go in reverse! Dijkhuizen and Koornwinder showed that if (\mathcal{A}, Δ) is a Hopf $*$ -algebra which is spanned by the matrix entries of its finite-dimensional unitary corepresentations, then there is a compact quantum group (A, Δ) such that \mathcal{A} is given by A .

Drinfeld's approach to quantum groups starts with a Lie group G , the Lie algebra \mathfrak{g} and the enveloping algebra $U(\mathfrak{g})$ which is naturally a Hopf algebra. It is this Hopf algebra which is deformed to get a "quantum group". If we start with $SU(2)$, and we now take the $*$ -representations of this deformed enveloping algebra, we naturally get a Hopf $*$ -algebra which is isomorphic to the Hopf $*$ -algebra associated to $C(SU_\mu(G))$.

So we are really studying the "dual" world to what is often understood by the term "quantum group".

More algebra

We can go in reverse! Dijkhuizen and Koornwinder showed that if (\mathcal{A}, Δ) is a Hopf $*$ -algebra which is spanned by the matrix entries of its finite-dimensional unitary corepresentations, then there is a compact quantum group (A, Δ) such that \mathcal{A} is given by A .

Drinfeld's approach to quantum groups starts with a Lie group G , the Lie algebra \mathfrak{g} and the enveloping algebra $U(\mathfrak{g})$ which is naturally a Hopf algebra. It is this Hopf algebra which is deformed to get a "quantum group". If we start with $SU(2)$, and we now take the $*$ -representations of this deformed enveloping algebra, we naturally get a Hopf $*$ -algebra which is isomorphic to the Hopf $*$ -algebra associated to $C(SU_\mu(G))$.

So we are really studying the "dual" world to what is often understood by the term "quantum group".

More algebra

We can go in reverse! Dijkhuizen and Koornwinder showed that if (\mathcal{A}, Δ) is a Hopf $*$ -algebra which is spanned by the matrix entries of its finite-dimensional unitary corepresentations, then there is a compact quantum group (A, Δ) such that \mathcal{A} is given by A .

Drinfeld's approach to quantum groups starts with a Lie group G , the Lie algebra \mathfrak{g} and the enveloping algebra $U(\mathfrak{g})$ which is naturally a Hopf algebra. It is this Hopf algebra which is deformed to get a “quantum group”. If we start with $SU(2)$, and we now take the $*$ -representations of this deformed enveloping algebra, we naturally get a Hopf $*$ -algebra which is isomorphic to the Hopf $*$ -algebra associated to $C(SU_\mu(G))$.

So we are really studying the “dual” world to what is often understood by the term “quantum group”.

More algebra

We can go in reverse! Dijkhuizen and Koornwinder showed that if (\mathcal{A}, Δ) is a Hopf $*$ -algebra which is spanned by the matrix entries of its finite-dimensional unitary corepresentations, then there is a compact quantum group (A, Δ) such that \mathcal{A} is given by A .

Drinfeld's approach to quantum groups starts with a Lie group G , the Lie algebra \mathfrak{g} and the enveloping algebra $U(\mathfrak{g})$ which is naturally a Hopf algebra. It is this Hopf algebra which is deformed to get a "quantum group". If we start with $SU(2)$, and we now take the $*$ -representations of this deformed enveloping algebra, we naturally get a Hopf $*$ -algebra which is isomorphic to the Hopf $*$ -algebra associated to $C(SU_\mu(G))$.

So we are really studying the "dual" world to what is often understood by the term "quantum group".

More algebra

We can go in reverse! Dijkhuizen and Koornwinder showed that if (\mathcal{A}, Δ) is a Hopf $*$ -algebra which is spanned by the matrix entries of its finite-dimensional unitary corepresentations, then there is a compact quantum group (A, Δ) such that \mathcal{A} is given by A .

Drinfeld's approach to quantum groups starts with a Lie group G , the Lie algebra \mathfrak{g} and the enveloping algebra $U(\mathfrak{g})$ which is naturally a Hopf algebra. It is this Hopf algebra which is deformed to get a "quantum group". If we start with $SU(2)$, and we now take the $*$ -representations of this deformed enveloping algebra, we naturally get a Hopf $*$ -algebra which is isomorphic to the Hopf $*$ -algebra associated to $C(SU_\mu(G))$.

So we are really studying the "dual" world to what is often understood by the term "quantum group".

More algebra

We can go in reverse! Dijkhuizen and Koornwinder showed that if (\mathcal{A}, Δ) is a Hopf $*$ -algebra which is spanned by the matrix entries of its finite-dimensional unitary corepresentations, then there is a compact quantum group (A, Δ) such that \mathcal{A} is given by A .

Drinfeld's approach to quantum groups starts with a Lie group G , the Lie algebra \mathfrak{g} and the enveloping algebra $U(\mathfrak{g})$ which is naturally a Hopf algebra. It is this Hopf algebra which is deformed to get a "quantum group". If we start with $SU(2)$, and we now take the $*$ -representations of this deformed enveloping algebra, we naturally get a Hopf $*$ -algebra which is isomorphic to the Hopf $*$ -algebra associated to $C(SU_\mu(G))$.

So we are really studying the "dual" world to what is often understood by the term "quantum group".

Back to analysis

- ▶ For a compact group G , we start with $C(G)$, and find a Haar measure to form $L^2(G)$.
- ▶ Then $C(G)$ is a C^* -algebra acting on $C(G)$.
- ▶ Consider the strong operator topology closure: this gives us $L^\infty(G)$, a von Neumann algebra.
- ▶ Δ extends to $L^\infty(G)$: it has the same formula.
- ▶ $L^\infty(G)$ has a predual: $L^1(G)$.
- ▶ Then Δ induces a Banach algebra product on $L^1(G)$, which is the usual convolution product.

Back to analysis

- ▶ For a compact group G , we start with $C(G)$, and find a Haar measure to form $L^2(G)$.
- ▶ Then $C(G)$ is a C^* -algebra acting on $C(G)$.
- ▶ Consider the strong operator topology closure: this gives us $L^\infty(G)$, a von Neumann algebra.
- ▶ Δ extends to $L^\infty(G)$: it has the same formula.
- ▶ $L^\infty(G)$ has a predual: $L^1(G)$.
- ▶ Then Δ induces a Banach algebra product on $L^1(G)$, which is the usual convolution product.

Back to analysis

- ▶ For a compact group G , we start with $C(G)$, and find a Haar measure to form $L^2(G)$.
- ▶ Then $C(G)$ is a C^* -algebra acting on $C(G)$.
- ▶ Consider the strong operator topology closure: this gives us $L^\infty(G)$, a von Neumann algebra.
- ▶ Δ extends to $L^\infty(G)$: it has the same formula.
- ▶ $L^\infty(G)$ has a predual: $L^1(G)$.
- ▶ Then Δ induces a Banach algebra product on $L^1(G)$, which is the usual convolution product.

Back to analysis

- ▶ For a compact group G , we start with $C(G)$, and find a Haar measure to form $L^2(G)$.
- ▶ Then $C(G)$ is a C^* -algebra acting on $L^2(G)$.
- ▶ Consider the strong operator topology closure: this gives us $L^\infty(G)$, a von Neumann algebra.
- ▶ Δ extends to $L^\infty(G)$: it has the same formula.
- ▶ $L^\infty(G)$ has a predual: $L^1(G)$.
- ▶ Then Δ induces a Banach algebra product on $L^1(G)$, which is the usual convolution product.

Back to analysis

- ▶ For a compact group G , we start with $C(G)$, and find a Haar measure to form $L^2(G)$.
- ▶ Then $C(G)$ is a C^* -algebra acting on $L^2(G)$.
- ▶ Consider the strong operator topology closure: this gives us $L^\infty(G)$, a von Neumann algebra.
- ▶ Δ extends to $L^\infty(G)$: it has the same formula.
- ▶ $L^\infty(G)$ has a predual: $L^1(G)$.
- ▶ Then Δ induces a Banach algebra product on $L^1(G)$, which is the usual convolution product.

Back to analysis

- ▶ For a compact group G , we start with $C(G)$, and find a Haar measure to form $L^2(G)$.
- ▶ Then $C(G)$ is a C^* -algebra acting on $L^2(G)$.
- ▶ Consider the strong operator topology closure: this gives us $L^\infty(G)$, a von Neumann algebra.
- ▶ Δ extends to $L^\infty(G)$: it has the same formula.
- ▶ $L^\infty(G)$ has a predual: $L^1(G)$.
- ▶ Then Δ induces a Banach algebra product on $L^1(G)$, which is the usual convolution product.

Interlude: KMS condition

- ▶ The Haar state on $C(G)$ or $C_r^*(\Gamma)$ is a *trace*:

$$\varphi(ab) = \varphi(ba) \quad (a, b \in A).$$

- ▶ For a compact quantum group, this is true if and only if we really have a Kac algebra.
- ▶ So not so for $C(SU_\mu(2))$ say.
- ▶ But φ is KMS.
- ▶ Loosely speaking, this means that there is an automorphism σ of \mathcal{A} such that

$$\varphi(ab) = \varphi(b\sigma(a)) \quad (a, b \in \mathcal{A}).$$

Interlude: KMS condition

- ▶ The Haar state on $C(G)$ or $C_r^*(\Gamma)$ is a *trace*:

$$\varphi(ab) = \varphi(ba) \quad (a, b \in A).$$

- ▶ For a compact quantum group, this is true if and only if we really have a Kac algebra.
- ▶ So not so for $C(SU_\mu(2))$ say.
- ▶ But φ is KMS.
- ▶ Loosely speaking, this means that there is an automorphism σ of \mathcal{A} such that

$$\varphi(ab) = \varphi(b\sigma(a)) \quad (a, b \in \mathcal{A}).$$

Interlude: KMS condition

- ▶ The Haar state on $C(G)$ or $C_r^*(\Gamma)$ is a *trace*:

$$\varphi(ab) = \varphi(ba) \quad (a, b \in A).$$

- ▶ For a compact quantum group, this is true if and only if we really have a Kac algebra.
- ▶ So not so for $C(SU_\mu(2))$ say.
- ▶ But φ is KMS.
- ▶ Loosely speaking, this means that there is an automorphism σ of \mathcal{A} such that

$$\varphi(ab) = \varphi(b\sigma(a)) \quad (a, b \in \mathcal{A}).$$

Interlude: KMS condition

- ▶ The Haar state on $C(G)$ or $C_r^*(\Gamma)$ is a *trace*:

$$\varphi(ab) = \varphi(ba) \quad (a, b \in A).$$

- ▶ For a compact quantum group, this is true if and only if we really have a Kac algebra.
- ▶ So not so for $C(SU_\mu(2))$ say.
- ▶ But φ is KMS.
- ▶ Loosely speaking, this means that there is an automorphism σ of \mathcal{A} such that

$$\varphi(ab) = \varphi(b\sigma(a)) \quad (a, b \in \mathcal{A}).$$

Interlude: KMS condition

- ▶ The Haar state on $C(G)$ or $C_r^*(\Gamma)$ is a *trace*:

$$\varphi(ab) = \varphi(ba) \quad (a, b \in A).$$

- ▶ For a compact quantum group, this is true if and only if we really have a Kac algebra.
- ▶ So not so for $C(SU_\mu(2))$ say.
- ▶ But φ is KMS.
- ▶ Loosely speaking, this means that there is an automorphism σ of \mathcal{A} such that

$$\varphi(ab) = \varphi(b\sigma(a)) \quad (a, b \in \mathcal{A}).$$

For compact quantum groups

Can do the same thing:

- $(A, \Delta) \implies$ Haar state $\varphi \implies$ Hilbert space H
- \implies von Neumann algebra M
- \implies predual M_* has structure of a Banach algebra.

If we start with discrete Γ then we get $A(\Gamma)$, the *Fourier algebra*: this encodes information about positive definite functions on Γ , and so forth.

If we start with $C(SU_\mu(2))$, we get: Who knows?

For compact quantum groups

Can do the same thing:

- $(A, \Delta) \implies$ Haar state $\varphi \implies$ Hilbert space H
- \implies von Neumann algebra M
- \implies predual M_* has structure of a Banach algebra.

If we start with discrete Γ then we get $A(\Gamma)$, the *Fourier algebra*:
this encodes information about positive definite functions on Γ ,
and so forth.

If we start with $C(SU_\mu(2))$, we get: Who knows?

For compact quantum groups

Can do the same thing:

- $(A, \Delta) \implies$ Haar state $\varphi \implies$ Hilbert space H
- \implies von Neumann algebra M
- \implies predual M_* has structure of a Banach algebra.

If we start with discrete Γ then we get $A(\Gamma)$, the *Fourier algebra*: this encodes information about positive definite functions on Γ , and so forth.

If we start with $C(SU_\mu(2))$, we get: Who knows?

For compact quantum groups

Can do the same thing:

- $(A, \Delta) \implies$ Haar state $\varphi \implies$ Hilbert space H
- \implies von Neumann algebra M
- \implies predual M_* has structure of a Banach algebra.

If we start with discrete Γ then we get $A(\Gamma)$, the *Fourier algebra*: this encodes information about positive definite functions on Γ , and so forth.

If we start with $C(SU_\mu(2))$, we get: *Who knows?*

For compact quantum groups

Can do the same thing:


- $(A, \Delta) \implies$ Haar state $\varphi \implies$ Hilbert space H
- \implies von Neumann algebra M
- \implies predual M_* has structure of a Banach algebra.

If we start with discrete Γ then we get $A(\Gamma)$, the *Fourier algebra*: this encodes information about positive definite functions on Γ , and so forth.

If we start with $C(SU_\mu(2))$, we get: Who knows?

Projectivity

If \mathfrak{A} is a Banach algebra, then in the category of left \mathfrak{A} -modules, P is *projective* if:




Complication: it might be impossible to solve this diagram problem for purely topological reasons. So we insist that the map $X \rightarrow Y$ is *admissible*, that is, there is a bounded linear (but not necessarily \mathfrak{A} -linear) right inverse.

Then \mathfrak{A} is (left) projective if, as a module over itself, it is projective.

Projectivity

If \mathfrak{A} is a Banach algebra, then in the category of left \mathfrak{A} -modules, P is *projective* if:




Complication: it might be impossible to solve this diagram problem for purely topological reasons. So we insist that the map $X \rightarrow Y$ is *admissible*, that is, there is a bounded linear (but not necessarily \mathfrak{A} -linear) right inverse.

Then \mathfrak{A} is (left) projective if, as a module over itself, it is projective.

Projectivity

If \mathfrak{A} is a Banach algebra, then in the category of left \mathfrak{A} -modules, P is *projective* if:




Complication: it might be impossible to solve this diagram problem for purely topological reasons. So we insist that the map $X \rightarrow Y$ is *admissible*, that is, there is a bounded linear (but not necessarily \mathfrak{A} -linear) right inverse.

Then \mathfrak{A} is (left) projective if, as a module over itself, it is projective.

Projectivity

If \mathfrak{A} is a Banach algebra, then in the category of left \mathfrak{A} -modules, P is *projective* if:



Complication: it might be impossible to solve this diagram problem for purely topological reasons. So we insist that the map $X \rightarrow Y$ is *admissible*, that is, there is a bounded linear (but not necessarily \mathfrak{A} -linear) right inverse.

Then \mathfrak{A} is (left) projective if, as a module over itself, it is projective.

For group algebras

- ▶ For a locally compact group G , we can still form $L^1(G)$ with convolution.
- ▶ Then $L^1(G)$ is projective if and only if G is compact.
- ▶ Similarly, the Fourier algebra $A(\Gamma)$ can be defined in general.
- ▶ If we work in the right category, then $A(\Gamma)$ is projective if and only if Γ is discrete.

The “right category” is the category of *completely bounded maps*.

For group algebras

- ▶ For a locally compact group G , we can still form $L^1(G)$ with convolution.
- ▶ Then $L^1(G)$ is projective if and only if G is compact.
- ▶ Similarly, the Fourier algebra $A(\Gamma)$ can be defined in general.
- ▶ If we work in the right category, then $A(\Gamma)$ is projective if and only if Γ is discrete.

The “right category” is the category of *completely bounded maps*.

For group algebras

- ▶ For a locally compact group G , we can still form $L^1(G)$ with convolution.
- ▶ Then $L^1(G)$ is projective if and only if G is compact.
- ▶ Similarly, the Fourier algebra $A(\Gamma)$ can be defined in general.
 - ▶ If we work in the right category, then $A(\Gamma)$ is projective if and only if Γ is discrete.

The “right category” is the category of *completely bounded maps*.

For group algebras

- ▶ For a locally compact group G , we can still form $L^1(G)$ with convolution.
- ▶ Then $L^1(G)$ is projective if and only if G is compact.
- ▶ Similarly, the Fourier algebra $A(\Gamma)$ can be defined in general.
- ▶ If we work in the right category, then $A(\Gamma)$ is projective if and only if Γ is discrete.

The “right category” is the category of *completely bounded maps*.

For group algebras

- ▶ For a locally compact group G , we can still form $L^1(G)$ with convolution.
- ▶ Then $L^1(G)$ is projective if and only if G is compact.
- ▶ Similarly, the Fourier algebra $A(\Gamma)$ can be defined in general.
- ▶ If we work in the right category, then $A(\Gamma)$ is projective if and only if Γ is discrete.

The “right category” is the category of *completely bounded maps*.

For compact quantum groups

Let M_* be the predual convolution algebra associated to a (locally) compact quantum group (A, Δ) . Again, we work with completely bounded maps.

- ▶ If M_* is projective, then A is compact.
- ▶ The converse is tricky!
- ▶ Abstract nonsense implies that M_* is projective if and only if

$$\Delta_* : M_* \widehat{\otimes} M_* \rightarrow M_*$$

the product map ($\widehat{\otimes}$ is the topological tensor product which linearises bilinear, completely bounded maps) has a completely bounded right inverse, which is an M_* module map.

- ▶ (Daws) If this inverse map is contractive, then A is a Kac algebra.
- ▶ The proof ends up showing that the modular automorphism of φ must be trivial.

For compact quantum groups

Let M_* be the predual convolution algebra associated to a (locally) compact quantum group (A, Δ) . Again, we work with completely bounded maps.

- ▶ If M_* is projective, then A is compact.
- ▶ The converse is tricky!
- ▶ Abstract nonsense implies that M_* is projective if and only if

$$\Delta_* : M_* \widehat{\otimes} M_* \rightarrow M_*$$

the product map ($\widehat{\otimes}$ is the topological tensor product which linearises bilinear, completely bounded maps) has a completely bounded right inverse, which is an M_* module map.

- ▶ (Daws) If this inverse map is contractive, then A is a Kac algebra.
- ▶ The proof ends up showing that the modular automorphism of φ must be trivial.

For compact quantum groups

Let M_* be the predual convolution algebra associated to a (locally) compact quantum group (A, Δ) . Again, we work with completely bounded maps.

- ▶ If M_* is projective, then A is compact.
- ▶ The converse is tricky!
- ▶ Abstract nonsense implies that M_* is projective if and only if

$$\Delta_* : M_* \widehat{\otimes} M_* \rightarrow M_*$$

the product map ($\widehat{\otimes}$ is the topological tensor product which linearises bilinear, completely bounded maps) has a completely bounded right inverse, which is an M_* module map.

- ▶ (Daws) If this inverse map is contractive, then A is a Kac algebra.
- ▶ The proof ends up showing that the modular automorphism of φ must be trivial.

For compact quantum groups

Let M_* be the predual convolution algebra associated to a (locally) compact quantum group (A, Δ) . Again, we work with completely bounded maps.

- ▶ If M_* is projective, then A is compact.
- ▶ The converse is tricky!
- ▶ Abstract nonsense implies that M_* is projective if and only if

$$\Delta_* : M_* \widehat{\otimes} M_* \rightarrow M_*$$

the product map ($\widehat{\otimes}$ is the topological tensor product which linearises bilinear, completely bounded maps) has a completely bounded right inverse, which is an M_* module map.

- ▶ (Daws) If this inverse map is contractive, then A is a Kac algebra.
- ▶ The proof ends up showing that the modular automorphism of φ must be trivial.

For compact quantum groups

Let M_* be the predual convolution algebra associated to a (locally) compact quantum group (A, Δ) . Again, we work with completely bounded maps.

- ▶ If M_* is projective, then A is compact.
- ▶ The converse is tricky!
- ▶ Abstract nonsense implies that M_* is projective if and only if

$$\Delta_* : M_* \widehat{\otimes} M_* \rightarrow M_*$$

the product map ($\widehat{\otimes}$ is the topological tensor product which linearises bilinear, completely bounded maps) has a completely bounded right inverse, which is an M_* module map.

- ▶ (Daws) If this inverse map is contractive, then A is a Kac algebra.
- ▶ The proof ends up showing that the modular automorphism of φ must be trivial.

For compact quantum groups

Let M_* be the predual convolution algebra associated to a (locally) compact quantum group (A, Δ) . Again, we work with completely bounded maps.

- ▶ If M_* is projective, then A is compact.
- ▶ The converse is tricky!
- ▶ Abstract nonsense implies that M_* is projective if and only if

$$\Delta_* : M_* \widehat{\otimes} M_* \rightarrow M_*$$

the product map ($\widehat{\otimes}$ is the topological tensor product which linearises bilinear, completely bounded maps) has a completely bounded right inverse, which is an M_* module map.

- ▶ (Daws) If this inverse map is contractive, then A is a Kac algebra.
- ▶ The proof ends up showing that the modular automorphism of φ must be trivial.

For compact quantum groups

Let M_* be the predual convolution algebra associated to a (locally) compact quantum group (A, Δ) . Again, we work with completely bounded maps.

- ▶ If M_* is projective, then A is compact.
- ▶ The converse is tricky!
- ▶ Abstract nonsense implies that M_* is projective if and only if

$$\Delta_* : M_* \widehat{\otimes} M_* \rightarrow M_*$$

the product map ($\widehat{\otimes}$ is the topological tensor product which linearises bilinear, completely bounded maps) has a completely bounded right inverse, which is an M_* module map.

- ▶ (Daws) If this inverse map is contractive, then A is a Kac algebra.
- ▶ The proof ends up showing that the modular automorphism of φ must be trivial.

For compact quantum groups

Let M_* be the predual convolution algebra associated to a (locally) compact quantum group (A, Δ) . Again, we work with completely bounded maps.

- ▶ If M_* is projective, then A is compact.
- ▶ The converse is tricky!
- ▶ Abstract nonsense implies that M_* is projective if and only if

$$\Delta_* : M_* \widehat{\otimes} M_* \rightarrow M_*$$

the product map ($\widehat{\otimes}$ is the topological tensor product which linearises bilinear, completely bounded maps) has a completely bounded right inverse, which is an M_* module map.

- ▶ (Daws) If this inverse map is contractive, then A is a Kac algebra.
- ▶ The proof ends up showing that the modular automorphism of φ must be trivial.